

Effects of stellarator transform on sawtooth oscillations in CTH

Jeffrey Herfindal



D.A. Ennis, J.D. Hanson, G.J. Hartwell, E.C. Howell, C.A. Johnson, S.F. Knowlton, X. Ma, D.A. Maurer, M.D. Pandya, N.A. Roberds, K.G. Ross, and P.J. Traverso



Understanding sawtooth physics and controlling their behavior is a critical tokamak research area



- First discovered in mid 1970s, physics understanding still an active area of research
- Large sawteeth have many deleterious effects on tokamak discharges:
 - Degradation of core confinement
 - Trigger for other MHD (ELMS, NTMS, locked modes) leading to disruption in some cases
- Control of large sawteeth is an important issue for ITER operation
- Small sawteeth can be beneficial by flushing impurities and helium ash from the core plasma

Sawteeth are periodic MHD initiated mixing events near the magnetic axis





- Mechanism responsible for the sawtooth crash is instability of the $m/n=1/1$ internal kink/tearing mode when $q_0 < 1$

Thermal energy transport leads to inverted sawteeth for $r > r_{inv}$



- Core plasma thermal energy inside the $q = 1$ inversion surface is rapidly transported and deposited outside of the inversion surface due to magnetic reconnection

Thermal energy transport leads to flat temperature profile after crash



- Study of non-ideal physics is important in order to understand the 1/1 mode evolution.

Various methods have been proposed for control of sawtooth period and amplitude

- Active control schemes based on different aspects of current understanding of $m/n=1/1$ mode stability physics
 - Energetic particle stabilization using ICRH or NBI
 - Changing local magnetic shear near the $q=1$ surface with ECCD and ECRH
 - Eliminate $q=1$ surface altogether by reversed shear operation

Various methods have been proposed for control of sawtooth period and amplitude

- Active control schemes based on different aspects of current understanding of $m/n=1/1$ mode stability physics
 - Energetic particle stabilization using ICRH or NBI
 - Changing local magnetic shear near the $q=1$ surface with ECCD and ECRH
 - Eliminate $q=1$ surface altogether by reversed shear operation
- 2D equilibrium shaping is known to effect sawtooth dynamics
 - High elongation is destabilizing¹
 - Triangularity is stabilizing²

1. Lütjens, H., Bondeson, A., Vlad G., Nucl. Fusion 32 (1992) 1625

2. Reimerdes, H. et al., Plasma Phys. Control Fusion 42 (2000)

Various methods have been proposed for control of sawtooth period and amplitude

- Active control schemes based on different aspects of current understanding of $m/n=1/1$ mode stability physics
 - Energetic particle stabilization using ICRH or NBI
 - Changing local magnetic shear near the $q=1$ surface with ECCD and ECRH
 - Eliminate $q=1$ surface altogether by reversed shear operation
- 2D equilibrium shaping is known to effect sawtooth dynamics
 - High elongation is destabilizing¹
 - Triangularity is stabilizing²
- Applied external $n=1$ fields on RFX-mod and DIII-D reduce the sawtooth period and amplitude³

1. Lütjens, H., Bondeson, A., Vlad G., Nucl. Fusion 32 (1992) 1625

2. Reimerdes, H. et al., Plasma Phys. Control Fusion 42 (2000)

3. Piron C. et al., Nucl. Fusion 56 (2016) 106012


Various methods have been proposed for control of sawtooth period and amplitude

- Active control schemes based on different aspects of current understanding of $m/n=1/1$ mode stability physics
 - Energetic particle stabilization using ICRH or NBI
 - Changing local magnetic shear near the $q=1$ surface with ECCD and ECRH
 - Eliminate $q=1$ surface altogether by reversed shear operation
- 2D equilibrium shaping is known to effect sawtooth dynamics
 - High elongation is destabilizing¹
 - Triangularity is stabilizing²

Can strong 3D shaping shed light on sawtooth physics and provide a passive control mechanism?

Compact Toroidal Hybrid (CTH) designed to study the effects of 3D shaping on MHD instabilities

- **Hybrid:** current driven within 3D equilibrium of a stellarator plasma
- Total rotational transform $t = t_{\text{vac}} + t_{\text{current}}$
 - t_{vac} from external stellarator coils (3D magnetic shaping)
 - t_{current} from plasma current
- CTH can vary the fractional transform, $t_{\text{vac}}(a)/t(a)$, from 4% to 100%



Major results

1. The observed sawtooth period and amplitude decrease with increasing 3D field
2. The sawtooth crash time does not change systematically with increasing 3D field
3. The decreasing sawtooth period and amplitude are correlated with increasing mean elongation
4. NIMROD resistive MHD simulations capture similar trend with sawtooth cycle period as seen in experiment



Outline

- Compact Toroidal Hybrid
- Sawtooth dynamics observed while varying the amount of 3D shaping
 1. Sawtooth period and amplitude change
 2. Sawtooth crash time
 3. Effect of mean elongation
 4. NIMROD simulations
- Summary

CTH: Flexible magnetic configuration in low aspect stellarator/tokamak hybrid



- **Helical Field coil** and **Toroidal Field coil** currents adjusted to modify vacuum rotational transform t_{vac}

$$R_0 = 0.75 \text{ m} \quad R/a \sim 4 \quad n_e \leq 5 \times 10^{19} \text{ m}^{-3} \quad T_e \leq 200 \text{ eV} \quad |B| \leq 0.7 \text{ T}$$




Ohmic coil allows induction of up to 95% of the total rotational transform from plasma current

- **Helical Field coil** and **Toroidal Field coil** currents adjusted to modify vacuum rotational transform t_{vac}
- **Central solenoid** drives $I_p \leq 80$ kA, dominating total transform



Sawtooth properties measured using a two-color SXR camera diagnostic




- The midplane SXR_M camera is used as an emissivity diagnostic to characterize sawtooth behavior with 3D shaping


Outline

- Compact Toroidal Hybrid
- Sawtooth dynamics observed while varying the amount of 3D shaping
 1. Sawtooth period and amplitude change
 2. Sawtooth crash time
 3. Effect of mean elongation
 4. NIMROD simulations
- Summary


Sawtooth oscillations observed on CTH exhibit behavior similar to that of axisymmetric tokamaks



Sawtooth oscillations observed on CTH exhibit behavior similar to that of axisymmetric tokamaks




Biorthogonal decomposition provides an empirical mode basis to characterize the sawteeth behavior




- Singular value decomposition (SVD) separates a spatio-temporal signal into two functions dependent on time or space: $f(x, t) = \sum_n A_n v_n(x) u_n(t)$

Reconstructed biorthogonal decomposition signals illustrate clear sawtoothing behavior and inversion radius





- Reconstructed SXR signals using the first two modes of BD
- Linear fit subtracted from each channel

Sawteeth observed in CTH exhibit similar scaling of inversion surface size as in tokamaks



- Normalized sawtooth inversion radius is proportional to $\tau(a) = 1/q(a)^1$

Observed inversion surface radius does not scale strongly with the amount of 3D shaping




- Normalized sawtooth inversion radius is proportional to $\tau(a) = 1/q(a)^1$


Outline

- Compact Toroidal Hybrid
- Two-color SXR camera system
- Sawtooth dynamics observed while varying the amount of 3D shaping
 1. Sawtooth period and amplitude change
 2. Sawtooth crash time
 3. Effect of mean elongation
 4. NIMROD simulations
- Summary



Increased 3D shaping observed to give rise to more frequent sawteeth




Sawtooth period and amplitude both decrease with application of higher 3D shaping



Sawtooth crash time appears to be unaffected by the amount of 3D shaping




1/1 mode dynamics observed throughout the sawtooth cycle



- The 1/1 mode spikes in amplitude during each sawtooth crash
- NIMROD CTH sawtooth simulations exhibit Kadomtsev-like sawtooth crashes

Large amplitude sawteeth not observed at high levels of 3D magnetic shaping



Outline


- Compact Toroidal Hybrid
- Two-color SXR camera system
- Sawtooth dynamics observed while varying the amount of 3D shaping
 1. Sawtooth period and amplitude change
 2. Sawtooth crash time
 3. Effect of mean elongation
 4. NIMROD simulations
- Summary

Elongation destabilizes $m/n = 1/1$ mode in tokamaks


- For axisymmetric plasmas $\kappa = b/a$
- To understand the possible effect of 3D elongation on our sawtooth observations we employ a mean elongation, κ , computed by VMEC¹
- This definition of κ reduces to the conventional definition of b/a if applied to an axisymmetric torus
- κ calculated on the last closed flux surface as a proxy for κ at the $q=1$ surface

1. ArchMiller et al., Phys. of Plasmas 21, 056113 (2014)

Shorter period sawteeth observed at higher levels of mean elongation



Decreased sawtooth amplitude also correlated with increasing mean elongation




Outline

- Compact Toroidal Hybrid
- Sawtooth dynamics observed while varying the amount of 3D shaping
 1. Sawtooth period and amplitude change
 2. Sawtooth crash time
 3. Mean elongation – comparison to 2D results
 4. NIMROD simulations
- Summary

Resistive MHD simulation of CTH sawteeth and $m/n = 1/1$ mode activity using NIMROD


2D
 $t_{\text{vac}} = 0$




- Axisymmetric plasma simulated with similar parameters to CTH

Fully 3D case exhibits a strongly shaped large amplitude 1/1 island and enhanced stochasticity


2D
 $t_{\text{vac}} = 0$



3D
 $t_{\text{vac}} = 0.033$




Linearly unstable MHD eigenfunction composed of a single $n=1$ mode in an axisymmetric configuration




For the axisymmetric case a single growing $n = 1$ mode is observed

Linearly unstable MHD eigenfunction has a rich toroidal harmonic content due helical shaping




For the axisymmetric case a single growing $n = 1$ mode is observed





With stellarator symmetry, the eigenfunction contains harmonics with $jN_{fp} \pm 1^1$

NIMROD simulations reproduce sawtooth cycling consistent with experiment



- Equilibrium represented by the Fourier numbers 0, 5, 10, ...

NIMROD simulations with higher levels of stellarator transform produce shorter period sawteeth



Summary

- The observed sawtooth period and amplitude decrease with increased 3D shaping using stellarator transform
- The sawtooth crash time is not strongly correlated with the amount of 3D stellarator field
- The decreased sawtooth period and amplitude are correlated with increasing edge vacuum rotational transform and mean elongation
- NIMROD resistive MHD simulations capture similar trend on the effect of 3D equilibrium shaping with sawtooth cycle period as seen in experiment