

Math 2650 - Linear Differential Equations

A. J. Meir

Copyright (C) A. J. Meir. All rights reserved.

This worksheet is for educational use only. No part of this publication may be reproduced or transmitted for profit in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system without prior written permission from the author. Not for profit distribution of the software is allowed without prior written permission, providing that the worksheet is not modified in any way and full credit to the author is acknowledged.

Separation of Variables

Separation of variables is an analytical method that may be used to solve certain d.e.s (and certain i.v.p.s). At the heart of the method is the chain rule.

Consider a (first order) differential equation that may be written as

$$\frac{dx}{dt} = f(x) g(t).$$

That is, the derivative of the function is equal to the product of a function of only x and a function of only t (the r.h.s. of the equation).

If $f(x) \neq 0$ we can write

$$\frac{1}{f(x)} \frac{dx}{dt} = g(t).$$

Recall $x = x(t)$, so if we can find a function $H(x)$ with $\frac{dH}{dx} = \frac{1}{f(x)}$, using the chain rule we have that

$$\frac{d}{dt}H(x) = \frac{d}{dx}H(x) \frac{dx}{dt} = \frac{1}{f(x)} \frac{dx}{dt}$$

so

$$\frac{d}{dt}H(x) = g(t).$$

Integrating both sides of the equation w.r.t. t we get

$$H(x) = \int g(t) dt + c$$

and since

$$H(x) = \int \frac{1}{f(x)} dx,$$

the solution of the d.e. is given (implicitly) by

$$\int \frac{1}{f(x)} dx = \int g(t) dt + C.$$

Note: the above is an implicit solution. Also note the arbitrary constant C , this is the general solution. If we are also given an initial condition we can use it to solve for C and obtain a particular solution.

Examples:

1.

$$\frac{dx}{dt} = kx$$

if $x \neq 0$, separating the variables

$$\frac{1}{x} dx = k dt$$

integrating

$$\ln(|x|) = kt + C$$

exponentiating

$$|x| = e^{kt + C}$$

so

$$|x| = C_1 e^{kt} \quad \text{where } C_1 = e^C$$

and absorbing the sign into the constant C_1 we get the solution

$$x(t) = C_2 e^{kt}$$

another solution is

$$x(t) = 0.$$

What is the interval of existence of these solutions?

Exercise: verify that the above are solutions of the differential equation.

2.

$$\frac{dx}{dt} = x^2 t$$

if $x \neq 0$

$$\frac{dx}{x^2} = t dt$$

a solution is given by

$$\int \frac{1}{x^2} dx = \int t dt + C$$

```
> restart;
> int(1/x^2, x);
```

$$-\frac{1}{x}$$

```
> int(t, t);
```

$$\frac{1}{2} t^2$$

So a solution is

$$-\frac{1}{x} = \frac{t^2}{2} + C.$$

Solving for x

[> **solve(-1/x = t^2/2+C,x);**

$$-\frac{2}{t^2 + 2C}$$

we obtain

$$x(t) = -\frac{2}{t^2 + 2C}$$

or

$$x(t) = \frac{2}{C - t^2}.$$

another solution is

$$x(t) = 0.$$

What is the interval of existence of these solutions? What does it depend on?

Exercise: verify that the above are solutions of the differential equation.

3.

$$\frac{dx}{dt} = x^2 \sin(t) + x \sin(t)$$

rewrite the equation as

$$\frac{dx}{dt} = (x^2 + x) \sin(t)$$

if $x^2 + x \neq 0$

$$\frac{dx}{x^2 + x} = \sin(t) dt$$

a solution is given by

$$\int \frac{1}{x^2 + x} dx = \int \sin(t) dt + C$$

[> **int(1/(x^2+x),x);**

$$\ln(x) - \ln(x + 1)$$

[> **int(sin(t),t);**

$$-\cos(t)$$

So a solution is

$$\ln(x) - \ln(x + 1) = -\cos(t) + C$$

[> **solve(ln(x)-ln(x+1) = -cos(t)+C,x);**

$$-\frac{e^{-\cos(t) + C}}{-1 + e^{-\cos(t) + C}}$$

so

$$x(t) = \frac{1}{C e^{\cos(t)} - 1}$$

[> **solve(x^2+x=0);**

0, -1

so additional solutions are

$$x(t) = 0 \quad \text{and} \quad x(t) = -1.$$

What is the interval of existence of these solutions? What does it depend on?

Exercise: verify that the above are solutions of the differential equation.

Problem:

Solve the equation

$$\frac{dx}{dt} = \frac{x}{t}.$$