

A. HORN'S RESULT ON MATRICES WITH PRESCRIBED SINGULAR VALUES AND EIGENVALUES*

TIN-YAU TAM[†]

Abstract. We give a new proof of a classical result of A. Horn on the existence of a matrix with prescribed singular values and eigenvalues.

Key words. Eigenvalues, singular values.

AMS subject classifications. 15A45, 15A18.

Let $A \in \mathbb{C}_{n \times n}$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A arranged in the order $|\lambda_1| \geq \dots \geq |\lambda_n|$. The singular values of A are the nonnegative square roots of the eigenvalues of the positive semi-definite matrix A^*A and are denoted by $s_1 \geq \dots \geq s_n$. Weyl's inequalities [7] provide a very nice relation between the eigenvalues and singular values of A :

$$\prod_{j=1}^k |\lambda_j| \leq \prod_{j=1}^k s_j, \quad k = 1, \dots, n-1, \quad (0.1)$$

$$\prod_{j=1}^n |\lambda_j| = \prod_{j=1}^n s_j. \quad (0.2)$$

The equality follows from two ways of expressing the absolute value of the determinant of A . A. Horn [2] established the converse of Weyl's result.

THEOREM 0.1. (A. Horn) If $|\lambda_1| \geq \dots \geq |\lambda_n|$ and $s_1 \geq \dots \geq s_n$ satisfy (0.1) and (0.2), then there exists $A \in \mathbb{C}_{n \times n}$ such that $\lambda_1, \dots, \lambda_n$ are the eigenvalues and s_1, \dots, s_n are the singular values of A .

Horn's original proof is divided into two cases: (i) $s_n \neq 0$ (the nonsingular case) and (ii) $s_n = 0$ (the singular case. There is a typo: $C_{m,m+1} = \gamma$ and $C_{i,i+1} = \alpha_i$ should be $C_{m+1,m} = \gamma$ and $C_{i+1,i} = \alpha_i$ on [2, p.6]). In this note we provide a new proof of Horn's result. Our proof differs from Horn's proof in two ways that (i) our proof is divided into two cases according to $\lambda_1 = 0$ and $\lambda_1 \neq 0$, and (ii) our induction technique is different. It is very much like Chan and Li's technique [1] (the same

*Received by the editors on Oct. 31, 2009. Accepted for publication on Month y, 200y. Handling Editor: Roger Horn.

[†]Department of Mathematics and Statistics, Auburn University, AL 36849-5310, USA (tamtiny@auburn.edu).

technique is also used in [8]) for proving another result of Horn [3] (on the diagonal entries and eigenvalues of a Hermitian matrix): ours is multiplicative and Chan and Li's is additive. See [4, Section 3.6] for a proof of Theorem 0.1 using the result of Horn [3]. Also see [5] for an extension of Weyl-Horn's result and a numerically stable construction of A . A technique similar to that of Chan and Li can be found in Thompson's earlier work [6] on the diagonal entries and singular values of a square matrix.

Proof. We divide the proof into two cases: nilpotent or not.

Case i: $\lambda_1 = 0$. Then $s_n = 0$ by (0.2) and we choose

$$A := \begin{pmatrix} 0 & s_1 & & \\ & 0 & \ddots & \\ & & & s_{n-1} \\ & & & 0 \end{pmatrix}$$

Case ii: $\lambda_1 \neq 0$. We will use induction on n . When $n = 2$, the matrix

$$A = \begin{pmatrix} \lambda_1 & \mu \\ 0 & \lambda_2 \end{pmatrix}$$

has singular values $s_1 \geq s_2$ if we set

$$\mu := (s_1^2 + s_2^2 - |\lambda_1|^2 - |\lambda_2|^2)^{1/2}.$$

Suppose that the statement of Theorem 0.1 is true for $\lambda_1 \neq 0$ when $n = m \geq 2$. Let $n = m + 1$ and let $j \geq 2$ be the largest index such that $s_{j-1} \geq |\lambda_1| \geq s_j$. Clearly $s_1 \geq \max\{|\lambda_1|, s_1 s_j / |\lambda_1|\} \geq \min\{|\lambda_1|, s_1 s_j / |\lambda_1|\}$. Then there exist 2×2 unitary matrices U_1 and V_1 such that

$$U_1 \begin{pmatrix} s_1 & \\ & s_j \end{pmatrix} V_1 = \begin{pmatrix} \lambda_1 & \mu' \\ 0 & s_1 s_j / |\lambda_1| \end{pmatrix},$$

where $\mu' = (s_1^2 + s_j^2 - |\lambda_1|^2 - s_1^2 s_j^2 / |\lambda_1|^2)^{1/2}$. Set $U_2 := U_1 \oplus I_{m-1}$, $V_2 := V_1 \oplus I_{m-1}$. Then

$$\begin{aligned} A_1 &:= U_2 \operatorname{diag}(s_1, s_j, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}) V_2 \\ &= \begin{pmatrix} \lambda_1 & \mu' \\ 0 & s_1 s_j / |\lambda_1| \end{pmatrix} \oplus \operatorname{diag}(s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}). \end{aligned}$$

It suffices to show that $(s_1 s_j / |\lambda_1|, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1})$ and $(\lambda_2, \dots, \lambda_{m+1})$ satisfy (0.1) and (0.2). Since $s_{j-1} \geq |\lambda_1| \geq |\lambda_2|$,

$$|\lambda_2| \leq \max\{s_1 s_j / |\lambda_1|, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}\}.$$

Moreover

$$\prod_{i=2}^k |\lambda_i| \leq |\lambda_1|^{k-1} \leq \prod_{i=2}^k s_i, \quad k = 2, \dots, j-1,$$

$$\prod_{i=2}^k |\lambda_i| = \frac{1}{|\lambda_1|} \prod_{i=1}^k |\lambda_i| \leq \frac{s_1 s_j}{|\lambda_1|} \prod_{i=2, i \neq j}^k s_i, \quad k = j, \dots, m, \quad \text{by (0.1)}$$

$$\prod_{i=2}^{m+1} |\lambda_i| = \frac{s_1 s_j}{|\lambda_1|} \prod_{i=2, i \neq j}^{m+1} s_i \quad \text{by (0.2)}.$$

We consider two cases: (a) $\lambda_2 = 0$ and apply Case i. (b) $\lambda_2 \neq 0$ and apply the induction hypothesis in Case ii. For both cases, there exist $m \times m$ unitary matrices U_3, V_3 such that

$$U_3 \operatorname{diag} \left(\frac{s_1 s_2}{|\lambda_1|}, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1} \right) V_3$$

is upper triangular with diagonal $(\lambda_2, \dots, \lambda_{m+1})$. Then $A = U_4 A_1 V_4$ is the desired matrix, where $U_4 := 1 \oplus U_3, V_4 := 1 \oplus V_3$. \square

Acknowledgement: The author is very thankful to Roger Horn for many valuable comments and suggestions.

REFERENCES

- [1] N.N. Chan and K.H. Li. Diagonal elements and eigenvalues of a real symmetric matrix. *J. Math. Anal. Appl.*, **91** (1983) 562–566.
- [2] A. Horn. On the eigenvalues of a matrix with prescribed singular values. *Proc. Amer. Math. Soc.*, **5** (1954) 4–7.
- [3] A. Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *Amer. J. Math.*, **76** (1954), 620–630.
- [4] R.A. Horn and C.R. Johnson. *Topics in Matrix Analysis*. Cambridge Univ. Press, 1991.
- [5] C.K. Li and R. Mathias, Construction of matrices with prescribed singular values and eigenvalues. *BIT* **41** (2001) 115–126.
- [6] R.C. Thompson. Singular values, diagonal elements, and convexity. *SIAM J. Appl. Math.*, **32** (1977) 39–63.
- [7] H. Weyl. Inequalities between the two kinds of eigenvalues of a linear transformation. *Proc. Nat. Acad. Sci. U. S. A.*, **35** (1949) 408–411.
- [8] H. Zha and Z. Zhang. A note on constructing a symmetric matrix with specified diagonal entries and eigenvalues. *BIT*, **35** (1995) 448–452.