

A FAMILY OF NONSEPARABLE SCALING FUNCTIONS AND SMOOTH COMPACTLY SUPPORTED TIGHT FRAMELETS

A. SAN ANTOLÍN AND R. A. ZALIK

ABSTRACT. Given integers b and d , with $d > 1$ and $|b| > 1$, we construct smooth nonseparable compactly supported refinable functions with dilation factor b that generate multiresolution analyses on $L^2(\mathbb{R}^d)$. These refinable functions are nonseparable, in the sense that they cannot be expressed as the product of two functions defined on lower dimensions. We use these scaling functions and a slight generalization of a theorem of Lai and Stöckler to construct smooth compactly supported tight framelets. Both the refinable functions and the framelets they generate can be made as smooth as desired. Estimates for the supports of these refinable functions and framelets, are given.

1. INTRODUCTION

Tight wavelet frames, or “framelets”, have recently become the focus of increased interest because they can be computed and applied just as easily as orthonormal wavelets, but are easier to construct. Moreover, since frames may have redundant terms, they are better suited for data transmission.

Han [12], and independently Ron and Shen [22], found necessary and sufficient conditions for translates and dilates of a set of functions to be a tight framelet. Ron and Shen also formulated what is known as the Unitary Extension Principle (UEP), which, in addition to its other applications, allowed Gröchenig and Ron to construct, for any dilation matrix, compactly supported framelets with any desired degree of smoothness (see [11]). Articles studying nonseparable frame or orthogonal wavelets include Ayache [1, 2, 3, 4], Karoui [14], Kovačević and Vetterli [16], Lai [18], Li [20], and Yang and Xue [29]. A closed form representation valid for any MRA orthonormal wavelet in $L^2(\mathbb{R}^d)$ was obtained by Zalik. It follows from [30, Theorem 3] and [31, Theorem 3 and Theorem 9] (there is a typographical error in [30, Theorem 3 and Theorem 4]: where it says *orthogonal matrix* should say *unitary matrix*). No such representation is currently known for tight framelets. In addition to the well-known work of Daubechies, the construction of compactly supported orthonormal wavelets was also studied by Lai [17] and Yang and Xue [28].

The UEP led to the Oblique Extension Principle (OEP), a method based on the UEP; it was developed by Chui, He and Stöckler ([7]), and independently by Daubechies, Han, Ron, and Shen ([9]), who gave the method its name. With the definition of the Fourier transform that we shall adopt in the next section, the OEP may be formulated as follows:

Key words and phrases. Fourier transform, multiresolution analysis, Riesz basis, scaling function, low pass filter, tight framelet, Paley–Wiener theorem for several complex variables.

2010 Mathematics Subject Classification: 42C40.

Theorem A. Let $B \in \mathbb{R}^{d \times d}$ be a dilation matrix preserving the lattice \mathbb{Z}^d . Let $\phi \in L^2(\mathbb{R}^d)$ be compactly supported and refinable, i.e.

$$\widehat{\phi}(B^* \mathbf{x}) = P(\mathbf{x}) \widehat{\phi}(\mathbf{x}),$$

where $P(\mathbf{x})$ is a trigonometric polynomial and B^* is the complex conjugate transpose of B . Assume moreover that $|\widehat{\phi}(\mathbf{0})| = 1$ and $|\widehat{\phi}(\mathbf{x})| \leq C(1+|\mathbf{x}|)^{-\alpha}$ for some $\alpha > d/2$. Let $S(\mathbf{x})$ be another trigonometric polynomial such that $S(\mathbf{x}) \geq \mathbf{0}$ and $S(\mathbf{0}) = 1$. Assume there are trigonometric polynomials or rational functions Q_ℓ , $\ell = 1, \dots, N$, that satisfy the OEP condition

$$(1) \quad S(B^* \mathbf{x}) P(\mathbf{x}) \overline{P(\mathbf{x} + \mathbf{j})} + \sum_{\ell=0}^N Q_\ell(\mathbf{x}) \overline{Q_\ell(\mathbf{x} + \mathbf{j})} \\ = \begin{cases} S(\mathbf{x}) & \text{if } \mathbf{j} = \mathbf{0}, \\ 0 & \text{if } \mathbf{j} \in (\mathbb{Z}^d / B^*(\mathbb{Z}^d)) \setminus B^*(\mathbb{Z}^d) \end{cases}.$$

If

$$\widehat{\psi}_\ell(\mathbf{x}) := Q_\ell(\mathbf{x}) \widehat{\phi}(\mathbf{x}), \quad \ell = 1, \dots, N,$$

then $\Psi = \{\psi_1, \dots, \psi_N\}$ is a tight framelet in $L^2(\mathbb{R}^d)$.

This version of the OEP is a straightforward consequence of [22, Corollary 6.7].

In [19], Lai and Stöckler used the OEP to provide two methods for constructing compactly supported framelets in $L^2(\mathbb{R}^d)$ for the dilation matrix $2I_{d \times d}$. The first method is for refinable functions whose masks P are quadrature mirror filters, whereas the second method is for masks that satisfy the so-called sub-QMF condition:

Theorem B. ([19, Theorem 3.4]) Let $B = 2I_{d \times d}$ and suppose that $P(\mathbf{x})$ satisfies the condition

$$\sum_{\gamma \in \{0, 1/2\}^d} |P(\mathbf{t} + \gamma)|^2 \leq 1,$$

let

$$\mathcal{P}(\mathbf{t}) := (P(\mathbf{t} + \gamma); \gamma \in \{0, 1/2\}^d)^T,$$

and let

$$\mathcal{M}(\mathbf{t}) := 2^{-d/2} \left(e^{i2\pi \delta \cdot (\mathbf{t} + \gamma)}; \gamma \in \{0, 1/2\}^d, \delta \in \{0, 1\}^d \right)$$

be the polyphase matrix, where γ denotes the row index and δ denotes the column index. Let the $2^d \times 1$ matrix function $G(\mathbf{t})$ be defined by

$$G(\mathbf{t}) := \mathcal{M}^*(\mathbf{t}) \mathcal{P}(\mathbf{t}) = (L_\delta(2\mathbf{t}); \delta \in \{0, 1\}^d)^T,$$

where $\mathcal{M}^*(\mathbf{t})$ denotes the complex conjugate transpose of $\mathcal{M}(\mathbf{t})$, and assume that there are nonnegative polynomials $\tilde{P}_j(\mathbf{t})$, $j = 1, \dots, M$, such that

$$\sum_{\delta \in \{0, 1\}^d} |L_\delta(\mathbf{t})|^2 + \sum_{j=1}^M [\tilde{P}_j(\mathbf{t})]^2 = 1.$$

Then there exist $2^d + M$ trigonometric polynomials Q_ℓ . $\ell = 1, \dots, 2d + M$ such that P and the polynomials Q_ℓ satisfy the OEP condition with $S(\mathbf{x}) = \mathbf{1}$.

Lai and Stöckler also gave a number of examples using multivariate box splines.

A generalization of their results to vector-valued frames was obtained by Charina, Chui and He [6].

In the present paper we use Theorem A and a generalization of Theorem B to construct another family of smooth tight framelets with compact support, for systems with dilation matrix $bI_{d \times d}$. For all the members of this family, the frame generators are nonseparable and their masks may be obtained by spectral factorization. Also, estimates for their supports and degrees of smoothness are given. We should also point out that all members of this family have the same number of generators for each given dimension d : $|b|^d + 2d$. Thus, in contrast to [23, 24], this number is independent of the smoothness of the frame generators. Since the number of generators for an orthonormal wavelet associated with an MRA with dilation matrix $bI_{d \times d}$ is $|b|^d - 1$, we see that $|b|^d + 2d$ is asymptotically close for large values of d .

The remainder of this paper is organized as follows. The next section summarizes the notation and definitions that will be used. Section 3 is devoted to the construction of compactly supported nonseparable smooth scaling functions in a b -MRA in $L^2(\mathbb{R}^d)$, $d > 1$. In Section 4 we construct smooth tight framelets of compactly supported functions with any dilation factor b , from the scaling functions defined in Section 3.

2. NOTATION AND DEFINITIONS

The sets of strictly positive integers, integers, and real numbers will be denoted by \mathbb{N} , \mathbb{Z} and \mathbb{R} respectively, and the set of complex numbers by \mathbb{C} . Given $c \in \mathbb{R}$ and $\mathbf{x} = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$, we define $c\mathbf{x} := (cx_1, cx_2, \dots, cx_d)$ and $c(\mathbb{Z}) := \{ck : k \in \mathbb{Z}\}$. Given $m \in \mathbb{N}$, we will also use the following notation: $\Delta_m := \{0, 1, \dots, m-1\}$, and $\Gamma_m := \{0, 1/m, \dots, (m-1)/m\}$. By Δ_b^d we mean the d -fold cartesian product of Δ_b with itself, and Γ_b^d is similarly defined; $\mathbf{t} = (t_1, \dots, t_d)$, $\mathbf{y} = (y_1, \dots, y_d)$, $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_d)$, and $\boldsymbol{\delta} = (\delta_1, \dots, \delta_d)$ will be vectors in \mathbb{R}^d , $\mathbf{z} = (z_1, \dots, z_d) = \mathbf{x} + i\mathbf{y} \in \mathbb{C}^d$, and $\mathbf{0}$ will denote the zero vector in \mathbb{R}^d . In what follows, b will always be an integer such that $m := |b| > 1$.

A function $\phi \in L^2(\mathbb{R}^d)$, $d > 1$ is said to be separable if there exist $a \in \mathbb{N}$, $1 \leq a < d$, and two functions $\varphi \in L^2(\mathbb{R}^a)$ and $\theta \in L^2(\mathbb{R}^{d-a})$ such that ϕ can be expressed as

$$(2) \quad \phi(x_1, \dots, x_a, x_{a+1}, \dots, x_d) = \varphi(x_1, \dots, x_a)\theta(x_{a+1}, \dots, x_d)$$

If $\phi \in L^2(\mathbb{R}^d)$ is not a separable function, then it is called a nonseparable function.

A sequence $\{\phi_n\}_{n=1}^\infty$ of elements in a separable Hilbert space \mathbb{H} is a *frame* for \mathbb{H} if there exist constants $C_1, C_2 > 0$ such that

$$C_1 \|h\|^2 \leq \sum_{n=1}^{\infty} |\langle h, \phi_n \rangle|^2 \leq C_2 \|h\|^2, \quad \forall h \in \mathbb{H},$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product on \mathbb{H} . The constants C_1 and C_2 are called *frame bounds*. The definition implies that a frame is a complete set of elements in \mathbb{H} . A frame $\{\phi_n\}_{n=1}^\infty$ is *tight* if we can choose $C_1 = C_2$.

If there are constants $C_1, C_2 > 0$ such that for any sequence $\{\alpha_{\mathbf{k}}\}_{\mathbf{k} \in \mathbb{Z}^d} \in l^2(\mathbb{Z}^n)$

$$C_1 \sum_{\mathbf{k} \in \mathbb{Z}^d} |\alpha_{\mathbf{k}}|^2 \leq \left\| \sum_{\mathbf{k} \in \mathbb{Z}^d} \alpha_{\mathbf{k}} \phi_{\mathbf{k}} \right\|^2 \leq C_2 \sum_{\mathbf{k} \in \mathbb{Z}^d} |\alpha_{\mathbf{k}}|^2,$$

then $\{\phi_n\}_{n=1}^\infty$ is called a *Riesz sequence*. It is a *Riesz basis* of the closure of its linear span.

Let $B \in \mathbb{R}^{d \times d}$ be a dilation matrix preserving the lattice \mathbb{Z}^d . A *multiresolution analysis* (MRA) in $L^2(\mathbb{R}^d)$ (generated by B) is a sequence $\{V_j; j \in \mathbb{Z}\}$ of closed linear subspaces of $L^2(\mathbb{R}^d)$ such that:

- (i) $V_j \subset V_{j+1}$ for every $j \in \mathbb{Z}$.
- (ii) For every $j \in \mathbb{Z}$, $f(\mathbf{t}) \in V_j$ if and only if $f(B\mathbf{t}) \in V_{j+1}$.
- (iii) $\bigcup_{j \in \mathbb{Z}} V_j$ is dense in $L^2(\mathbb{R}^d)$.
- (iv) There exists a function $\phi \in V_0$, called a *scaling function*, such that $\{\phi(\mathbf{x} - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz basis for V_0 .

Let $B \in \mathbb{R}^{d \times d}$ be a dilation matrix preserving the lattice \mathbb{Z}^d . A set of functions $\Psi = \{\psi_1, \dots, \psi_n\} \subset L^2(\mathbb{R}^d)$ is called a wavelet frame, if the system

$$\{|\det B|^{j/2} \psi_\ell(B^j \mathbf{t} + \mathbf{k}); j \in \mathbb{Z}, \mathbf{k} \in \mathbb{Z}^d, 1 \leq \ell \leq n\}$$

is a frame for $L^2(\mathbb{R}^d)$. If this system is a tight frame for $L^2(\mathbb{R}^d)$ then Ψ is called a tight wavelet frame. In particular, a tight wavelet frame with frame constant equal to 1 is called a *tight framelet*.

Let $b \in \mathbb{Z}$ and $|b| > 1$. By an MRA with dilation factor b , or a b -MRA, in $L^2(\mathbb{R}^d)$, we mean a multiresolution analysis with dilation matrix $bI_{d \times d}$, where $I_{d \times d}$ denotes the identity matrix on \mathbb{R}^d .

The simplest way to construct an MRA in $L^2(\mathbb{R}^d)$ is by multiplying scaling functions of lower dimensions: assume that φ is a scaling function for a b -MRA in $L^2(\mathbb{R}^a)$, $1 \leq a < d$, and that θ is a scaling function for a b -MRA in $L^2(\mathbb{R}^{d-a})$. Then the function ϕ defined as in (2) is the scaling function of a b -MRA in $L^2(\mathbb{R}^d)$.

If ϕ is a scaling function of an MRA in $L^2(\mathbb{R}^d)$, then conditions (i), (ii) and (iv) imply that ϕ is *refinable*, i.e.,

$$(3) \quad \widehat{\phi}(B^* \mathbf{t}) = P(\mathbf{t}) \widehat{\phi}(\mathbf{t}) \quad \text{a.e. on } \mathbb{R}^d$$

where B^* is the complex conjugate transpose of B , and the *low pass filter* $P(\mathbf{t})$ is a \mathbb{Z}^d -periodic essentially bounded function.

Conversely, if ϕ is a refinable function and we define

$$(4) \quad V_j := \overline{\text{span}}\{|\det B|^{j/2} \phi(B^j \mathbf{t} - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}, \quad j \in \mathbb{Z}$$

we see that conditions (i) and (ii) in the definition of MRA are satisfied. Thus, we say that a refinable function $\phi \in L^2(\mathbb{R}^d)$ *generates an MRA* if $\{\phi(\mathbf{t} - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz sequence and the subspaces V_j satisfy condition (iii).

Let \widehat{f} denote the Fourier transform of the function f . Thus, if $f \in L^1(\mathbb{R}^d)$, $\mathbf{t}, \mathbf{x} \in \mathbb{R}^d$,

$$\widehat{f}(\mathbf{x}) := \int_{\mathbb{R}^d} f(\mathbf{t}) e^{-2\pi i \mathbf{t} \cdot \mathbf{x}} d\mathbf{t},$$

where $\mathbf{t} \cdot \mathbf{x}$ denotes the usual inner product of vectors \mathbf{t} and \mathbf{x} .

3. CONSTRUCTION OF SMOOTH NONSEPARABLE SCALING FUNCTIONS

In this section we construct smooth nonseparable compactly supported scaling functions of a b -MRA in $L^2(\mathbb{R}^d)$, $d > 1$. We begin with

Proposition 1. *Let $m \in \mathbb{N}$ and*

$$(5) \quad p(t) := \frac{1}{m} \sum_{\delta \in \Delta_m} e^{-2\pi i \delta t} = \frac{1}{m} \frac{e^{-2\pi i m t} - 1}{e^{-2\pi i t} - 1},$$

let $n_1, n_2 \in \mathbb{N}$ with $n_1 < n_2$, and let the trigonometric polynomial $P(\mathbf{t})$ in \mathbb{R}^d be defined by

$$(6) \quad P(\mathbf{t}) := \frac{1-i}{2} \left(\prod_{j=1}^d |p(t_j)|^{2n_1} + i \prod_{j=1}^d |p(t_j)|^{2n_2} \right).$$

Then

$$(7) \quad \sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 = \frac{1}{2} \prod_{j=1}^d \left(\sum_{\gamma \in \Gamma_m} |p(t_j + \gamma)|^{4n_1} \right) + \frac{1}{2} \prod_{j=1}^d \left(\sum_{\gamma \in \Gamma_m} |p(t_j + \gamma)|^{4n_2} \right) \leq 1.$$

Moreover,

$$\sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 < 1,$$

except for a countable set of points.

Proof. Let $\gamma = (\gamma_1, \dots, \gamma_d)$ and $\tilde{\gamma} = (\gamma_2, \dots, \gamma_d)$. We have

$$\begin{aligned} \sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 &= \sum_{\gamma \in \Gamma_m^d} \frac{1}{2} \left(\prod_{j=1}^d |p(t_j + \gamma_j)|^{4n_1} + \prod_{j=1}^d |p(t_j + \gamma_j)|^{4n_2} \right) \\ &= \sum_{\gamma \in \Gamma_m} |p(t_1 + \gamma)|^{4n_1} \left(\sum_{\tilde{\gamma} \in \Gamma_m^{d-1}} \frac{1}{2} \left(\prod_{j=2}^d |p(t_j + \gamma_j)|^{4n_1} \right) \right) \\ &\quad + \sum_{\gamma \in \Gamma_m} |p(t_1 + \gamma)|^{4n_2} \left(\sum_{\tilde{\gamma} \in \Gamma_m^{d-1}} \frac{1}{2} \left(\prod_{j=2}^d |p(t_j + \gamma_j)|^{4n_2} \right) \right). \end{aligned}$$

Repeating this procedure a finite number of times we obtain the desired identity in (7), and the inequality follows from (5).

Finally note that (5) also implies that $p(t) = 1$ if and only if $t \in \mathbb{Z}$. This implies that the left-hand side of (7) is strictly less than 1 if and only if $\mathbf{t} \notin \mathbb{Z}^d$. \square

The following statement may be found in ([27, Appendix A.2]). The proof is straightforward and will be omitted.

Lemma C. *Let C^0 be the class of continuous functions in $L^2(\mathbb{R}^d)$, and let C^r , $r = 1, 2, \dots$ be the class of functions f such that all partial derivatives of f of order not greater than r are continuous and in $L^2(\mathbb{R}^d)$. If*

$$|\widehat{f}(\mathbf{t})| \leq C(1 + |\mathbf{t}|)^{-N-\varepsilon}$$

for some integer $N \geq d$ and $\varepsilon > 0$, then f is in C^{N-d} .

We now prove.

Proposition 2. *Let $b \in \mathbb{Z}$, assume that $m := |b| > 1$, and let P be defined by (6) with $n_1 < n_2$. Then*

$$(8) \quad \prod_{j=1}^{\infty} P(b^{-j}\mathbf{t})$$

converges uniformly on compact subsets of \mathbb{R}^d , and the function ϕ whose Fourier transform $\widehat{\phi}$ is defined by (8) has the following properties: it is continuous and square-integrable on \mathbb{R}^d ; if $2n_1 > d + 1$ it is in continuity class C^{2n_1-d-1} ; it has support in $[-n_2, n_2]^d$. Moreover, $\widehat{\phi}(\mathbf{0}) = 1$ and $\{ \phi(\cdot - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d \}$ is a Riesz sequence.

Proof. Let S be a compact subset of \mathbb{R}^d . Clearly P satisfies a Lipschitz condition on S ; thus

$$|P(\mathbf{t}_1) - P(\mathbf{t}_2)| \leq C|\mathbf{t}_1 - \mathbf{t}_2|.$$

Since (5) implies that $p(0) = 1$, we conclude from (6) that $P(\mathbf{0}) = 1$. Therefore, the preceding displayed inequality implies that

$$|P(\mathbf{t}) - 1| \leq C|\mathbf{t}|,$$

whence

$$|P(b^{-j}\mathbf{t}) - 1| \leq m^{-j}C|\mathbf{t}|.$$

This implies that the infinite product converges uniformly on compact subsets of \mathbb{R}^d .

To see that $\widehat{\phi}(\mathbf{0}) = 1$, it suffices to note that $\prod_{j=1}^n P(b^{-j}\mathbf{0}) = 1$ for all n .

Proceeding as above, we also conclude that

$$\prod_{j=1}^{\infty} p(m^{-j}t)$$

converges uniformly on compact subsets of \mathbb{R} to a function $A(t)$. Let $c(t)$ denote the characteristic function of the set $[0, 1]$. Since

$$[0, 1) = \bigcup_{k=0}^{m-1} \left[\frac{k}{m}, \frac{k+1}{m} \right),$$

we readily see that

$$\widehat{c}(mt) = p(t)\widehat{c}(t),$$

and we conclude that

$$A(t) = \widehat{c}(t) = e^{-\pi it} \frac{\sin \pi t}{\pi t}.$$

Therefore, since $|p(t)|^2$ is an even function,

$$(9) \quad \prod_{j=1}^{\infty} |p(b^{-j}t)|^{2n} = \prod_{j=1}^{\infty} |p(m^{-j}t)|^{2n} = \left| \frac{\sin \pi t}{\pi t} \right|^{2n}.$$

Since

$$|P(\mathbf{t})| \leq \prod_{j=1}^d |p(t_j)|^{2n_1},$$

we have

$$(10) \quad |\widehat{\phi}(\mathbf{t})| \leq \prod_{j=1}^d \left(\frac{\sin \pi t_j}{\pi t_j} \right)^{2n_1} \leq K(1 + |\mathbf{t}|)^{-2n_1}.$$

Thus $\widehat{\phi} \in L^2(\mathbb{R}^d)$, and therefore $\phi(\mathbf{t})$ is well defined as the inverse Fourier transform of $\widehat{\phi}$. Moreover, if $2n_1 > d + 1$, Lemma C implies that ϕ is in continuity class C^{2n_1-d-1} .

Let

$$v(\mathbf{t}) := \prod_{j=1}^d c(t_j),$$

$$V(\mathbf{t}) := \sum_{\mathbf{k} \in \mathbb{Z}^d} |\widehat{v}(\mathbf{t} + \mathbf{k})|^2,$$

and

$$\Phi(\mathbf{t}) := \sum_{\mathbf{k} \in \mathbb{Z}^d} |\widehat{\phi}(\mathbf{t} + \mathbf{k})|^2.$$

Since $\{d(\cdot - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is an orthonormal sequence, we know that $V(\mathbf{t}) = 1$ a.e.; but $\Phi(\mathbf{t}) \leq V(\mathbf{t})$. Thus, $\Phi(\mathbf{t}) \leq 1$ a.e. (This also follows from [25, p. 113]).

On the other hand,

$$|P(\mathbf{t})| \geq \prod_{j=1}^d |p(t_j)|^{2n_2},$$

and from (9) we see that

$$(11) \quad |\widehat{\phi}(\mathbf{t})| \geq \prod_{j=1}^d \left(\frac{\sin \pi t_j}{\pi t_j} \right)^{2n_2}.$$

Therefore,

$$\Phi(\mathbf{t}) \geq |\widehat{\phi}(\mathbf{t})|^2 \geq C > 0, \quad \text{a.e. in } [-\frac{1}{2}, \frac{1}{2}]^d.$$

Since $\Phi(\mathbf{t})$ is \mathbb{Z}^d -periodic, the preceding inequality holds on \mathbb{Z}^d , and we have established that $\{\phi(\cdot - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz sequence. It remains to prove that the support of ϕ is in $[-n_2, n_2]^d$. We will use an argument of Wojtaszczyk [27, p. 79].

The definition of $p(t)$ in Proposition 1 implies that

$$|p(t)|^2 = \frac{1}{m^2} \left(\sum_{l=0}^{m-1} e^{-2\pi i l t} \right) \left(\sum_{j=0}^{m-1} e^{2\pi i j t} \right) = \sum_{k=-(m-1)}^{m-1} a_k e^{-2\pi i k t}, \quad a_k \in \mathbb{R}.$$

This implies that if $n \in \mathbb{N}$, then

$$|p(t)|^{2n} = \sum_{k=-n(m-1)}^{n(m-1)} c_k e^{-2\pi i k t}, \quad c_k \in \mathbb{R},$$

which readily yields

$$P(\mathbf{t}) = \sum_{\mathbf{k} \in \Omega} \alpha_{\mathbf{k}} e^{-2\pi i \mathbf{k} \mathbf{t}}, \quad \alpha_{\mathbf{k}} \in \mathbb{C},$$

where $\Omega = [-n_2(m-1), n_2(m-1)]^d \cap \mathbb{Z}^d$.

Given $N \in \mathbb{N}$, let

$$\Pi_N(\mathbf{t}) = \prod_{j=1}^N P(b^{-j}\mathbf{t}),$$

and observe that Π_N may be written as

$$\Pi_N(\mathbf{t}) = \sum_{\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(N)}; \mathbf{k}^{(j)} \in \Omega} \alpha_{\mathbf{k}^{(1)}} \cdots \alpha_{\mathbf{k}^{(N)}} e^{2\pi i \mathbf{t} \cdot (\sum_{j=1}^N b^{-j} \mathbf{k}^{(j)})}.$$

Since $\mathbf{k}^{(j)} = (k_1^{(j)}, \dots, k_d^{(j)}) \in \mathbb{Z}^d$ with $|k_s^{(j)}| \leq n_2(m-1)$, $s = 1, \dots, d$, then $0 \leq \sum_{j=1}^N |b^{-j} k_s^{(j)}| \leq 2n_2(m-1) \sum_{j=1}^{\infty} m^{-j} = n_2$. Hence,

$$(12) \quad \sum_{j=1}^N b^{-j} \mathbf{k}^{(j)} \in [-n_2, n_2]^d.$$

Assume now that f is a function in the Schwartz class of \mathbb{R}^d such that $\widehat{f}(\mathbf{t}) = 0$ for every \mathbf{t} in $[-n_2, n_2]^d$. Bearing in mind that $|\Pi_N(\mathbf{t})| \leq 1$ for every \mathbf{t} in \mathbb{R}^d and using the Lebesgue Dominated Convergence Theorem, we obtain:

$$\begin{aligned} \int_{\mathbb{R}^d} \phi(\mathbf{t}) \widehat{f}(\mathbf{t}) d\mathbf{t} &= \int_{\mathbb{R}^d} \widehat{\phi}(\mathbf{t}) f(-\mathbf{t}) d\mathbf{t} = \lim_{N \rightarrow \infty} \int_{\mathbb{R}^d} \Pi_N(\mathbf{t}) f(-\mathbf{t}) d\mathbf{t} \\ &= \lim_{N \rightarrow \infty} \sum_{\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(N)}; \mathbf{k}^{(j)} \in \Omega} \alpha_{\mathbf{k}^{(1)}} \cdots \alpha_{\mathbf{k}^{(N)}} \int_{\mathbb{R}^d} e^{-2\pi i \mathbf{t} \cdot (\sum_{j=1}^N b^{-j} \mathbf{k}^{(j)})} f(-\mathbf{t}) d\mathbf{t} \\ &= \lim_{N \rightarrow \infty} \sum_{\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(N)}; \mathbf{k}^{(j)} \in \Omega} \alpha_{\mathbf{k}^{(1)}} \cdots \alpha_{\mathbf{k}^{(N)}} \widehat{f}\left(-\sum_{j=1}^N m^{-j} \mathbf{k}^{(j)}\right) = 0, \end{aligned}$$

and we conclude that the support of ϕ is in $[-n_2, n_2]^d$. \square

Our estimate for the support of ϕ may also be obtained from (12) using the Paley–Wiener Theorem for several complex variables. ([26, Theorem 4.9]).

From, e.g. [15, Theorem J], we obtain the following multivariate version of a result that is well-known for the single variable and dyadic dilation case ([13, p. 46]):

Proposition D. *Let V_j , $j \in \mathbb{Z}$, be a sequence of closed subspaces in $L^2(\mathbb{R}^d)$ satisfying conditions (i), (ii) and (iv) of the definition of b -MRA. Assume moreover that the scaling function ϕ is such that the origin is a point of continuity of $|\widehat{\phi}|$. Then $\overline{\bigcup_{j \in \mathbb{Z}^n} V_j} = L^2(\mathbb{R}^d)$ if and only if $\widehat{\phi}(\mathbf{0}) \neq 0$.*

We thus have:

Proposition 3. *The function ϕ , defined in Proposition 1, generates a b -MRA with low pass filter P .*

Proof. Proposition 2 implies that $\{\phi(\cdot - \mathbf{k}) : \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz sequence and that ϕ and the trigonometric polynomial P satisfy the refinement equation (3) with $A = bI_{d \times d}$. This implies that $V_j \subset V_{j+1}$, for every $j \in \mathbb{Z}$, where the subspaces V_j are defined by (4) with $A = bI_{d \times d}$. It remains to prove that $\bigcup_{j \in \mathbb{Z}} V_j$ is dense in $L^2(\mathbb{R}^d)$. This follows from Proposition 1 and Proposition D. \square

Proposition 4. *Let P be defined by (6) with $m \in \mathbb{N}$ and $n_1 < n_2$. Then there do not exist an integer a , $0 < a < d$, and two functions $f(\mathbf{t}_1)$ and $g(\mathbf{t}_2)$ defined on $L^2(\mathbb{R}^a)$ and $L^2(\mathbb{R}^{d-a})$ respectively and continuous at the origin, such that $f(\mathbf{t}_1)$ is \mathbb{Z}^a -periodic, $g(\mathbf{t}_2)$ is \mathbb{Z}^{d-a} -periodic, and*

$$P(t_1, \dots, t_a, t_{a+1}, \dots, t_d) = f(t_1, \dots, t_a)g(t_{a+1}, \dots, t_d) \quad \text{for } (t_1, \dots, t_d) \in \mathbb{R}^d.$$

Proof. We will argue by contradiction. Assume that there is an integer a , $1 \leq a < d$, and two functions $f(\mathbf{t})$ and $g(\mathbf{t})$ such as described in the statement of the proposition. Then, for $\mathbf{t} = (t_1, \dots, t_d) \in \mathbb{R}^d$,

$$(13) \quad \prod_{s=1}^d |p(t_s)|^{4n_1} + \prod_{s=1}^d |p(t_s)|^{4n_2} = 2f(t_1, \dots, t_a)g(t_{a+1}, \dots, t_d).$$

If $(t_1, \dots, t_a) = (t_1, \dots, t_1)$ and $(t_{a+1}, \dots, t_d) = (t_d, \dots, t_d)$, where $t_1, t_d \in \mathbb{R}$, we have

$$|p(t_1)|^{4n_1 a} |p(t_d)|^{4n_1(d-a)} + |p(t_1)|^{4n_2 a} |p(t_d)|^{4n_2(d-a)} = 2f(t_1, \dots, t_1)g(t_d, \dots, t_d).$$

In particular, if $t_d = 0$

$$(14) \quad |p(t_1)|^{4n_1 a} + |p(t_1)|^{4n_2 a} = 2f(t_1, \dots, t_1)g(0, \dots, 0).$$

Observe that $g(0, \dots, 0) \neq 0$ because $1 = f(0, \dots, 0)g(0, \dots, 0)$. Thus, substituting (14) into (13),

$$g(0, \dots, 0) \left(|p(t_1)|^{4n_1 a} |p(t_d)|^{4n_1(d-a)} + |p(t_1)|^{4n_2 a} |p(t_d)|^{4n_2(d-a)} \right) = 2 \left(|p(t_1)|^{4n_1 a} + |p(t_1)|^{4n_2 a} \right) g(t_d, \dots, t_d).$$

Therefore

$$(15) \quad |p(t_1)|^{4n_1 a} \left(g(0, \dots, 0) |p(t_d)|^{4n_1(d-a)} - 2g(t_d, \dots, t_d) \right) = |p(t_1)|^{4n_2 a} \left(g(0, \dots, 0) |p(t_d)|^{4n_2(d-a)} - 2g(t_d, \dots, t_d) \right).$$

Since $|p|^{4n_1}$ and $|p|^{4n_2}$ are different continuous functions, then at least one of the sets

$$S_1 := \{ t \in \mathbb{R} : g(0, \dots, 0) |p(t)|^{4n_1(d-a)} - 2g(t, \dots, t) \neq 0 \}$$

or

$$S_2 := \{ t \in \mathbb{R} : g(0, \dots, 0) |p(t)|^{4n_2(d-a)} - 2g(t, \dots, t) \neq 0 \}$$

must be nonempty. Without essential loss of generality we may assume that S_1 is nonempty. Define $S_p := \{ t \in [-1/2, 1/2]; p(t) \neq 0 \}$, and observe that S_p may differ from $[-1/2, 1/2]$ in an at most finite number of points.

Thus, if $t_d \in S_1$ and $t_1 \in S_p$, $t_1 \neq t_d$, (15) yields

$$(16) \quad -|p(t_1)|^{4(n_1-n_2)a} = \frac{g(0, \dots, 0) |p(t_d)|^{4n_2(d-a)} - 2g(t_d, \dots, t_d)}{g(0, \dots, 0) |p(t_d)|^{4n_1(d-a)} - 2g(t_d, \dots, t_d)}.$$

Since the variables in (16) are separated, we conclude that there is a constant C such that

$$|p(t_1)|^{4(n_1-n_2)a} = C$$

In fact, $C = 1$ because

$$\lim_{t \rightarrow 0, t \in S_p} |p(t)|^{4(n_1-n_2)a} = |p(0)|^{4(n_1-n_2)a} = 1.$$

We have thus reached a contradiction. \square

Combining Propositions 2, 3, and 4 we obtain

Theorem 1. *Let $b \in \mathbb{Z}$, assume that $m := |b| > 1$, let P be defined by (6) with $n_1 < n_2$, and let $\phi(\mathbf{t})$ be the function whose Fourier transform is given by (8). Then $\{\phi(\cdot - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz sequence, and ϕ has the following properties: it is a continuous, square-integrable, nonseparable scaling function of a b -MRA with lowpass filter $P(\mathbf{t})$, and its support is in $[-n_2, n_2]^d$. Moreover, if $2n_1 > d + 1$, then ϕ is in continuity class C^{2n_1-d-1} .*

Proof. The only thing that needs to be established is that ϕ is nonseparable: all the other statements follow directly from Propositions 2 and 3. We proceed by contradiction. Assume there exist $a \in \mathbb{N}$, $1 \leq a < d$, and two functions $\varphi \in L^2(\mathbb{R}^a)$ and $\theta \in L^2(\mathbb{R}^{d-a})$ such that

$$(17) \quad \phi(x_1, \dots, x_a, x_{a+1}, \dots, x_d) = \varphi(x_1, \dots, x_a)\theta(x_{a+1}, \dots, x_d) \quad \text{a.e.}$$

Since ϕ is compactly supported, also φ and θ must be compactly supported. Thus $\widehat{\phi}$, $\widehat{\varphi}$ and $\widehat{\theta}$ are continuous, and (17) implies that

$$\widehat{\phi}(t_1, \dots, t_a, t_{a+1}, \dots, t_d) = \widehat{\varphi}(t_1, \dots, t_a)\widehat{\theta}(t_{a+1}, \dots, t_d).$$

Since $1 = \widehat{\phi}(\mathbf{0}) = \widehat{\varphi}(0, \dots, 0)\widehat{\theta}(0, \dots, 0)$, then

$$\widehat{\varphi}(t_1, \dots, t_a) = [\widehat{\theta}(\mathbf{0})]^{-1}\widehat{\phi}(t_1, \dots, t_a, 0, \dots, 0)$$

and

$$\widehat{\theta}(t_{a+1}, \dots, t_d) = [\widehat{\varphi}(\mathbf{0})]^{-1}\widehat{\phi}(0, \dots, 0, t_{a+1}, \dots, t_d).$$

Observe that $P(t_1, \dots, t_a, 0, \dots, 0)$ and $P(0, \dots, 0, t_{a+1}, \dots, t_d)$ are continuous functions. Thus,

$$\begin{aligned} & P(t_1, \dots, t_a, t_{a+1}, \dots, t_d)\widehat{\phi}(t_1, \dots, t_a, t_{a+1}, \dots, t_d) \\ &= \widehat{\phi}(bt_1, \dots, bt_a, bt_{a+1}, \dots, bt_d) \\ &= \widehat{\varphi}(bt_1, \dots, bt_a)\widehat{\theta}(bt_{a+1}, \dots, bt_d) \\ &= P(t_1, \dots, t_a, 0, \dots, 0)P(0, \dots, 0, t_{a+1}, \dots, t_d)\widehat{\varphi}(t_1, \dots, t_a)\widehat{\theta}(t_{a+1}, \dots, t_d). \end{aligned}$$

But (11) implies that $\widehat{\phi}(\mathbf{t}) \neq 0$ a.e. By continuity, we therefore conclude that

$$P(t_1, \dots, t_a, t_{a+1}, \dots, t_d) = P(t_1, \dots, t_a, 0, \dots, 0)P(0, \dots, 0, t_{a+1}, \dots, t_d),$$

in contradiction of Proposition 4. \square

4. CONSTRUCTION OF COMPACTLY SUPPORTED TIGHT FRAMELETS

Note that the property that $\{\phi(\cdot - \mathbf{k}); \mathbf{k} \in \mathbb{Z}^d\}$ is a Riesz sequence is not used in this section.

Using the scaling functions that we obtained in the previous section, we now construct tight framelets $\Psi = \{\psi_1, \dots, \psi_n\}$ in $L^2(\mathbb{R}^d)$, $d > 1$, with any dilation factor b , such that functions ψ_ℓ , $\ell = 1, \dots, n$, are smooth and compactly supported. For this purpose we use a generalization of Theorem B to any dilation factor b . The proof is identical, and we include it for the sake of completeness. We have also included in the statement, a generalization of the algorithm implicit in the proof of Theorem B.

Theorem 2. Let $b \in \mathbb{Z}$, assume that $m := |b| > 1$, let $P(\mathbf{t})$ be a trigonometric polynomial defined on \mathbb{R}^d that satisfies the condition

$$\sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 \leq 1,$$

let

$$\mathcal{P}(\mathbf{t}) := (P(\mathbf{t} + \gamma); \gamma \in \Gamma_m^d)^T,$$

and let

$$(18) \quad \mathcal{M}(\mathbf{t}) := m^{-d/2} \left(e^{i2\pi\delta \cdot (\mathbf{t} + \gamma)}; \gamma \in \Gamma_m^d, \delta \in \Delta_m^d \right)$$

be the polyphase matrix, where γ denotes the row index and δ denotes the column index. Let the $m^d \times 1$ matrix function $G(\mathbf{t})$ be defined by

$$(19) \quad G(\mathbf{t}) := \mathcal{M}^*(\mathbf{t})\mathcal{P}(\mathbf{t}) = (L_\delta(b\mathbf{t}); \delta \in \Delta_m^d)^T,$$

where $\mathcal{M}^*(\mathbf{t})$ denotes the complex conjugate transpose of $\mathcal{M}(\mathbf{t})$. Suppose that there exist trigonometric polynomials $\tilde{P}_1, \dots, \tilde{P}_M$ such that

$$(20) \quad \sum_{\delta \in \Delta_m^d} |L_\delta(\mathbf{t})|^2 + \sum_{j=1}^M |\tilde{P}_j(\mathbf{t})|^2 = 1.$$

Let $N := m^d + M$ and let the $N \times 1$ matrix function $\mathcal{G}(\mathbf{t})$ be defined by

$$\mathcal{G}(\mathbf{t}) := \left(L_\delta(b\mathbf{t}); \delta \in \Delta_m^d, \tilde{P}_j(b\mathbf{t}); 1 \leq j \leq M \right)^T,$$

and

$$\tilde{\mathcal{Q}}(\mathbf{t}) := I_{N \times N} - \mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t}).$$

Let $H(\mathbf{t})$ denote the first $m^d \times N$ block matrix of $\tilde{\mathcal{Q}}(\mathbf{t})$,

$$\mathcal{Q}(\mathbf{t}) := \mathcal{M}(\mathbf{t})H(\mathbf{t}),$$

and let $[Q_1(\mathbf{t}), \dots, Q_N(\mathbf{t})]$ denote the first row of $\mathcal{Q}(\mathbf{t})$. Then the trigonometric polynomials P and Q_ℓ , $\ell = 1, \dots, N$ satisfy the identity (1) with $S(\mathbf{x}) = \mathbf{1}$.

Proof. Using (20), we have $\mathcal{G}^*(\mathbf{t})\mathcal{G}(\mathbf{t}) = 1$, then

$$\begin{aligned} \tilde{\mathcal{Q}}(\mathbf{t})\tilde{\mathcal{Q}}^*(\mathbf{t}) &= I_{N \times N} - 2\mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t}) + \mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t})\mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t}) \\ &= I_{N \times N} - \mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t}) = \tilde{\mathcal{Q}}(\mathbf{t}). \end{aligned}$$

Thus

$$\mathcal{G}(\mathbf{t})\mathcal{G}^*(\mathbf{t}) + \tilde{\mathcal{Q}}(\mathbf{t})\tilde{\mathcal{Q}}^*(\mathbf{t}) = I_{N \times N}.$$

Restricting to the first principal $m^d \times m^d$ blocks in the above matrices, we have

$$(21) \quad G(\mathbf{t})G^*(\mathbf{t}) + H(\mathbf{t})H^*(\mathbf{t}) = I_{m^d \times m^d}.$$

Since $\mathcal{M}(\mathbf{t})$ is unitary and by (21),

$$\mathcal{M}(\mathbf{t})G(\mathbf{t})G^*(\mathbf{t})\mathcal{M}^*(\mathbf{t}) + \mathcal{M}(\mathbf{t})H(\mathbf{t})H^*(\mathbf{t})\mathcal{M}^*(\mathbf{t}) = I_{m^d \times m^d}.$$

Thus,

$$\mathcal{P}(\mathbf{t})\mathcal{P}^*(\mathbf{t}) + \mathcal{Q}(\mathbf{t})\mathcal{Q}^*(\mathbf{t}) = I_{m^d \times m^d}$$

which is equivalent to saying that $P(\mathbf{t})$ and $Q_1(\mathbf{t}), \dots, Q_N(\mathbf{t})$ satisfy the equality (1) with $S(\mathbf{x}) = 1$ and with dilation matrix $B = bI_{d \times d}$. \square

We now need an auxiliary proposition.

Let h and v be trigonometric polynomials on \mathbb{R} such that

$$|h(bt)|^2 = 1 - \sum_{\gamma \in \Gamma_m} |p(t + \gamma)|^{4n_1} \quad \text{and} \quad |v(bt)|^2 = 1 - \sum_{\gamma \in \Gamma_m} |p(t + \gamma)|^{4n_2}.$$

To see that these polynomials exist note that, for example,

$$1 - \sum_{\gamma \in \Gamma_m} |p(t + \gamma)|^{4n_1} \geq 0,$$

and the assertion follows applying a lemma of Riesz (cf., e.g., [8, Lemma 6.1.3]), [21, Lemma 10, p. 102]). A similar argument shows that there exist two trigonometric polynomials u and w on \mathbb{R} , such that

$$|u(bt)|^2 = \sum_{\gamma \in \Gamma_m} |p(t + \gamma)|^{4n_1} \quad \text{and} \quad |w(bt)|^2 = \sum_{\gamma \in \Gamma_m} |p(t + \gamma)|^{4n_2}.$$

The coefficients of the polynomials h , v , u and w may be obtained by *spectral factorization* ([10]).

We have

Proposition 5. *Let $b \in \mathbb{Z}$, assume that $m := |b| > 1$, let p , n_1 , n_2 and P be as in Proposition 1, let the trigonometric polynomials h , v , u and w be as described in the preceding paragraph, let the trigonometric polynomials \tilde{P}_j be defined by*

$$(22) \quad \begin{aligned} \tilde{P}_j(\mathbf{t}) &:= \frac{\sqrt{2}}{2} h(t_j) \prod_{s=j+1}^d u(t_s), \quad j = 1, \dots, d-1, \\ \tilde{P}_d(\mathbf{t}) &:= \frac{\sqrt{2}}{2} h(t_d) \\ \tilde{P}_{d+j}(\mathbf{t}) &:= \frac{\sqrt{2}}{2} v(t_j) \prod_{s=j+1}^d w(t_s), \quad j = 1, \dots, d-1, \\ \tilde{P}_{2d}(\mathbf{t}) &:= \frac{\sqrt{2}}{2} v(t_d), \end{aligned}$$

and let the trigonometric polynomials $L_\delta(\mathbf{bt})$, $\delta \in \Delta_m^d$ be defined by (19). Then

$$\sum_{\delta \in \Delta_m^d} |L_\delta(\mathbf{t})|^2 + \sum_{j=1}^{2d} |\tilde{P}_j(\mathbf{t})|^2 = 1.$$

Proof. By Proposition 1 we know that

$$\sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 < 1,$$

except for a countable set of points. We also have

$$\begin{aligned}
& \sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 + \sum_{j=1}^{2d} |\tilde{P}_j(\mathbf{bt})|^2 \\
= & \frac{1}{2} \prod_{s=1}^d \left(\sum_{\gamma \in \Gamma_m} |p(t_s + \gamma)|^{4n_1} \right) + \frac{1}{2} \prod_{s=1}^d \left(\sum_{\gamma \in \Gamma_m} |p(t_s + \gamma)|^{4n_2} \right) \\
& + \frac{1}{2} \sum_{j=1}^{d-1} |h(bt_j)|^2 \prod_{s=j}^d |u(bt_s)|^2 + \frac{1}{2} |h(bt_d)|^2 \\
& + \frac{1}{2} \sum_{j=1}^{d-1} |v(bt_j)|^2 \prod_{s=j}^d |w(bt_s)|^2 + \frac{1}{2} |v(bt_d)|^2 \\
= & \frac{1}{2} \prod_{s=1}^d |u(bt_s)|^2 + \frac{1}{2} \prod_{s=1}^d |w(bt_s)|^2 \\
& + \frac{1}{2} |h(bt_1)|^2 \prod_{s=1}^d |u(bt_s)|^2 + \frac{1}{2} \sum_{j=2}^{d-1} |h(bt_j)|^2 \prod_{s=j}^d |u(bt_s)|^2 + \frac{1}{2} |h(bt_d)|^2 \\
& + \frac{1}{2} |v(bt_1)|^2 \prod_{s=1}^d |w(bt_s)|^2 + \frac{1}{2} \sum_{j=2}^{d-1} |v(bt_j)|^2 \prod_{s=j}^d |w(bt_s)|^2 + \frac{1}{2} |v(bt_d)|^2 \\
= & \frac{1}{2} (|u(bt_1)|^2 + |h(bt_1)|^2) \prod_{s=2}^d |u(bt_s)|^2 + \frac{1}{2} (|w(bt_1)|^2 + |v(bt_1)|^2) \prod_{s=2}^d |w(bt_s)|^2 \\
& + \frac{1}{2} \sum_{j=2}^{d-1} |h(bt_j)|^2 \prod_{s=j+1}^d |u(bt_s)|^2 + \frac{1}{2} |h(bt_d)|^2 \\
& + \frac{1}{2} \sum_{j=2}^{d-1} |v(bt_j)|^2 \prod_{s=j+1}^d |w(bt_s)|^2 + \frac{1}{2} |v(bt_d)|^2 \\
= & \frac{1}{2} \prod_{s=2}^d |u(bt_s)|^2 + \frac{1}{2} \prod_{s=2}^d |w(bt_s)|^2 + \frac{1}{2} \sum_{j=2}^{d-1} |h(bt_j)|^2 \prod_{s=j+1}^d |u(bt_s)|^2 + \frac{1}{2} |h(bt_d)|^2 \\
& + \frac{1}{2} \sum_{j=2}^{d-1} |v(bt_j)|^2 \prod_{s=j+1}^d |w(bt_s)|^2 + \frac{1}{2} |v(bt_d)|^2.
\end{aligned}$$

Repeating this procedure a finite number of times, we finally obtain

$$\begin{aligned}
& \sum_{(\rho_1, \dots, \rho_d) \in \Gamma_m^d} |P(t_1 + \rho_1, \dots, t_d + \rho_d)|^2 + \sum_{j=1}^{2d} |\tilde{P}(bt_1, \dots, bt_d)|^2 \\
= & \frac{1}{2} |u(bt_d)|^2 + \frac{1}{2} |w(bt_d)|^2 + \frac{1}{2} |h(bt_d)|^2 + \frac{1}{2} |v(bt_d)|^2 = 1,
\end{aligned}$$

i.e.,

$$\sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2 + \sum_{j=1}^{2d} |\tilde{P}_j(b\mathbf{t})|^2 = 1.$$

Finally, note that, since $\mathcal{M}(\mathbf{t})$ defined by (18) is unitary,

$$\sum_{\delta \in \Delta_m^d} |L_\delta(b\mathbf{t})|^2 = \mathcal{P}^*(\mathbf{t})\mathcal{M}(\mathbf{t})\mathcal{M}^*(\mathbf{t})\mathcal{P}(\mathbf{t}) = \sum_{\gamma \in \Gamma_m^d} |P(\mathbf{t} + \gamma)|^2.$$

□

The following theorem describes the construction of a tight smooth framelet of compact support for any dilation factor b .

Theorem 3. *Let $b \in \mathbb{Z}$ be such that $m := |b| > 1$, let $n_1, n_2 \in \mathbb{N}$ with $n_1 < n_2$ and $2n_1 > d + 1$, let $N := m^d + 2d$, and let the trigonometric polynomials $Q_1(\mathbf{t}), \dots, Q_N(\mathbf{t})$ be obtained by the algorithm described in Theorem 2 with $P(\mathbf{t})$ defined by (6) and the trigonometric polynomials $\tilde{P}_j(\mathbf{t})$, $j = 1, \dots, 2d$ defined as in Proposition 5. If*

$$\widehat{\phi}(\mathbf{t}) := \prod_{j=1}^{\infty} P(b^{-j}\mathbf{t}),$$

$$\widehat{\psi}_\ell(\mathbf{t}) := Q_\ell(\mathbf{t})\widehat{\phi}(\mathbf{t}), \quad \ell = 1, \dots, N.$$

and $\Psi = \{\psi_1, \dots, \psi_N\}$ is the set of inverse Fourier transforms of the functions $\widehat{\psi}_\ell$ defined in the preceding displayed identity, then Ψ is a tight framelet in $L^2(\mathbb{R}^d)$ with dilation factor b , and the functions $\psi_\ell(\mathbf{t})$ are square-integrable on \mathbb{R}^d , have support in $[-mn_2, mn_2]^d$, and are in continuity class C^{2n_1-d-1} .

Proof. That $\Psi = \{\psi_1, \dots, \psi_N\}$ is a tight framelet follows from Proposition 5, Theorem 2 and Theorem A.

Since the functions Q_ℓ are trigonometric polynomials and therefore bounded on \mathbb{R}^d , the smoothness of the functions ψ_ℓ follows from (10) and Lemma C.

If

$$I_n := [-(m-1)n_2, (m-1)n_2]^d \cap \mathbb{Z}^d,$$

then (6) and (1) imply that the polynomials Q_ℓ have representations of the form

$$Q_\ell(\mathbf{t}) = \sum_{\mathbf{k} \in I_n} \alpha_{\mathbf{k}} e^{-2\pi i \mathbf{k} \cdot \mathbf{t}}.$$

This implies that

$$\widehat{\psi}_\ell(\mathbf{t}) = \sum_{\mathbf{k} \in I_n} \alpha_{\mathbf{k}} \phi(\mathbf{t} - \mathbf{k}),$$

and since the support of ϕ is in $[-n_2, n_2]^d$, we conclude that the supports of the functions $\psi_\ell(\mathbf{t})$ are in $[-mn_2, mn_2]^d$. □

REFERENCES

- [1] A. Ayache, Bases multivariées d'ondelettes orthonormales à support compact et de régularité arbitraire, Ph.D. thesis, Ceremade, University of Paris Dauphine, 1997.
- [2] A. Ayache, Construction de bases orthonormées de $L^2(\mathbb{R}^2)$ non-séparables, à support compact et de régularité arbitrairement grande, C.R. Acad. Sci Paris, 325 Série 1 (1997) 17–20.
- [3] A. Ayache, Construction of non separable dyadic compactly supported orthonormal wavelet bases for $L^2(\mathbb{R}^2)$ of arbitrarily high regularity, Rev. Mat. Iberoamericana 15, No.1 (1999) 37–58.
- [4] A. Ayache, Some methods for constructing nonseparable, orthonormal, compactly supported wavelet bases, Appl. Comput. Harmon. Anal. 10 (2001) 99–111.
- [5] E. Belogay, Y. Wang, Arbitrarily smooth orthogonal non-separable wavelets in R^2 , SIAM J. Math. Anal. 30 (1999) 678–697.
- [6] M. Charina, C.K. Chui, W. He, Tight frames of compactly supported multivariate multi-wavelets, J. Comput. Appl. Math. 233 (2010) 2044–2061.
- [7] C.K. Chui, W. He, J. Stöckler, Compactly supported tight and sibling frames with maximum vanishing moments, Appl. Comput. Harmonic Anal. 13 (2002) 224–262.
- [8] I. Daubechies, *Ten lectures on wavelets*, SIAM, Philadelphia, 1992.
- [9] I. Daubechies, B. Han, A. Ron, Z. W. Shen, Framelets: MRA-based constructions of wavelet frames, Appl. Comput. Harmon. Anal. 14 (2003), 1–46.
- [10] T.N.T. Goodman, C.A. Micchelli, G. Rodriguez, S. Seatzu, Spectral factorization of Laurent polynomials, Adv. Comput. Math. 7 (1997), 429–454.
- [11] K. Gröchenig, A. Ron, Tight compactly supported wavelet frames of arbitrarily high smoothness, Proc. Amer. Math. Soc. 126 (1998) 1101–1107.
- [12] B. Han, On dual wavelet tight frames, Applied Comput. Harmon. Anal. 4 (1997) 380–413.
- [13] E. Hernández, G. Weiss; *A first course on Wavelets*, CRC Press Inc., 1996.
- [14] A. Karoui, A note on the construction of nonseparable wavelet bases and multiwavelet matrix filters of $L^2(\mathbb{R}^n)$, where $n \geq 2$, Electron. Res. Announc. of the Amer. Math. Soc. 9 (2003), 32–39.
- [15] K. S. Kazarian, A. San Antolin, Characterization of scaling functions in a frame multiresolution analysis in H_G^2 , L. De Carli, K. Kazarian and M. Milman (Eds.), Topics in classical analysis and applications in honor of Daniel Waterman, 118–140, World Sci. Publ., Hackensack, NJ, 2008.
- [16] E. Kovačević, M. Vetterli, Nonseparable multidimensional perfect reconstruction filter banks and wavelet bases for \mathcal{R}^n , IEEE Trans. Inform. Theory 38, no.2 (1992) 533–555.
- [17] M.-J. Lai, Construction of multivariate compactly supported orthonormal wavelets, Adv. Comput. Math. 25 (2006) 41–56.
- [18] M.-J. Lai, Methods for constructing nonseparable compactly supported orthonormal wavelets, in: D.X. Zhou (Ed.), Wavelet Analysis: Twenty Years Development, World Scientific, Singapore, 2002, pp. 231–251.
- [19] M.-J. Lai, J. Stöckler, Construction of multivariate compactly supported tight wavelet frames, Appl. Comput. Harmon. Anal. 21 (2006) 324–348.
- [20] Y.-Z Li, On the construction of a class of bidimensional nonseparable compactly supported wavelets, Proc. Amer. Math. Soc. 138 (2005) 3505–3513.
- [21] Y. Meyer, *Ondelettes et opérateurs. I*, Hermann, Paris, 1996 [English Translation: Wavelets and operators, Cambridge University Press, 1992]
- [22] A. Ron, Z.W. Shen, Affine systems in $L^2(\mathbb{R}^d)$: The analysis of the analysis operator, J. Funct. Anal. 148 (1997) 408–447.
- [23] A. Ron, Z.W. Shen, Compactly supported tight affine frames in $L^2(\mathbb{R}^d)$, Math. Comp. 67 (1998) 191–207.
- [24] A. Ron, Z.W. Shen, Construction of compactly supported affine frames in $L^2(\mathbb{R}^d)$, in: K.S. Lau (Ed.), Advances in Wavelets, Springer-Verlag, New York, 1998, pp. 27–49.
- [25] A. San Antolín, Characterization of low pass filters in a multiresolution analysis, Studia Math. 190(2) (2009), 99–116.
- [26] E. Stein, G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, Princeton, NJ 1971.
- [27] P. Wojtaszczyk, *A mathematical introduction to wavelets*, London Math. Soc., Student Texts 37, 1997.

- [28] Shouzhi Yang, Yanmei Xue, Construction of compactly supported nonseparable orthogonal wavelets of $L^2(\mathbb{R}^n)$. Bull. Iranian Math. Soc. 36 (2010) 183–199.
- [29] Y. Shouzhi, X. Yanmei, Construction of compactly supported conjugate symmetric complex tight wavelet frames, Int. J. Wavelets Multiresolut. Inf. Process. 8 (2010) 861–874.
- [30] R. A. Zalik, Representation of Orthonormal Multivariate Wavelets, in G. Chen and M–J. Lai, Eds. Splines and Wavelets: Athens 2005, Nashboro Press, Brentwood, TN 2005, pp. 507–515.
- [31] R. A. Zalik, Bases of translates and multiresolution analyses, Appl. Comput. Harmon. Anal. 24 (2008), no. 1, 41–57. Corrigendum, Appl. Comput. Harmon. Anal. 29 (2010), 121.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE ALICANTE, 03080 ALICANTE, SPAIN.

E-mail address: `angel.sanantolin@ua.es`

DEPARTMENT OF MATHEMATICS AND STATISTICS, AUBURN UNIVERSITY, AUBURN, AL. 36849–5310

E-mail address: `zalik@auburn.edu`