




NORTH
CAROLINA
**MUSEUM
of
NATURAL
SCIENCES**

**FOURTH COLLOQUIUM
ON CONSERVATION
OF MAMMALS
IN THE SOUTHEASTERN
UNITED STATES**

Edited by BRIAN R. CHAPMAN
and JOSHUA LAERM



Occasional Papers of the
North Carolina Museum of Natural Sciences and the
North Carolina Biological Survey
Number 12
Fall, 2000

GASTROINTESTINAL HELMINTH PARASITES OF BATS IN ALABAMA

Clayton D. Hilton and Troy L. Best

*Department of Biological Sciences and Alabama Agricultural Experiment Station,
331 Funchess Hall, Auburn University, AL 36849-5414*

Present address of CDH: Montgomery Zoo, P.O. Box ZEBRA, Montgomery, AL 36109

Gastrointestinal tracts of 324 bats representing 10 species from 27 locations in Alabama were examined. Helminth parasites occurred in 121 (37.3%) of the bats; 14 (4.3%) had cestodes, 47 (14.5%) had nematodes, and 77 (23.8%) had trematodes. Fifteen bats (4.6%) were parasitized by two types of helminths, and one had all three types of helminths. There was no significant difference between numbers of big brown bats (*Eptesicus fuscus*) and Brazilian free-tailed bats (*Tadarida brasiliensis*) that were parasitized, but *E. fuscus* was more likely to have trematodes and *T. brasiliensis* was more likely to have nematodes. Prevalences of parasites (percentage of infected hosts) were: *E. fuscus*, 26.6% ($n = 128$); red bat (*Lasiurus borealis*), 80.0% ($n = 10$); evening bat (*Nycticeius humeralis*), 40.0% ($n = 15$); southeastern myotis (*Myotis austroriparius*), 72.7% ($n = 11$); eastern pipistrelle (*Pipistrellus subflavus*), 71.1% ($n = 45$); *T. brasiliensis*, 25.7% ($n = 109$).

Bats are hosts to many internal parasites (Stiles and Nolan, 1931; Webster, 1973). Ubelaker's (1970) review of endoparasites of bats included one genus of acanthocephalan, 10 genera of cestodes, 24 genera of nematodes, 13 genera of protozoans, and 42 genera of trematodes. Endoparasites have been recorded from carnivorous (Peterson and Kirmse, 1969), frugivorous (Bray, 1984; Ubelaker et al., 1977), insectivorous (Duszynski et al., 1988), piscivorous (Fischthal and Martin, 1978; Zdzitowiecki and Rutkowska, 1980), and sanguivorous (Greenhall and Schmidt, 1988) bats. Despite the long scientific history documenting parasites in bats from throughout the world (Webster, 1973), endoparasites of bats in North America are poorly known (Coggins, 1988). There are only five published reports on internal parasites of bats from the southeastern United States; three are on helminth parasites of bats from Louisiana, Mississippi, Tennessee (Byrd and Macy, 1942), Texas, Louisiana (Martin, 1976), and Florida (Loftin, 1961), and two are on protozoan parasites of bats from Alabama (Wheat, 1975) and Florida (Foster, 1979).

There have been three parasitologic studies of bats in Alabama. White (1959) reported 30 species of acarine mites from 10 species of bats. Wheat (1975) described *Eimeria macyi* (Protozoa: Eimeriidae) from a *P. subflavus* that was collected in Lion's Den Cave, Clarke Co. More recently, Durden et al. (1992) recorded seven species of acarine mites on cohabitating *E. fuscus* and *T. brasiliensis* from Auburn University, Lee Co. None of these studies

have involved helminth parasites of bats. Because limited emphasis has been placed on helminthiasis of bats in any geographic area (Nickel and Hansen, 1967), we conducted this study to identify species of bats in Alabama that have a parasitic helminthofauna. Additional goals of this study were to examine variations in helminth faunas with respect to species of host, collection locality, sex of host, season, and interrelationships of parasitic helminths.

MATERIALS AND METHODS

A total of 324 specimens representing 10 of the 16 species of bats that occur in Alabama (Best et al., 1993; E. R. Hall, 1981) was examined from 27 locations during this study (Appendix I). Of these specimens, 88 were *E. fuscus* and 105 were *T. brasiliensis* taken from the attic of Samford Hall, Auburn University. Samford Hall is a large, four-story building with a double-roofed design that serves as a year-round residence for both species of bats (Durden et al., 1992; Henry et al., 2000). During each month from February through November 1990, ≤ 12 *E. fuscus* and *T. brasiliensis* were either collected from their roost sites by hand or trapped with a harp trap. Care was taken not to collect young-of-the-year or females that were caring for young. Other species of bats were collected February 1990-April 1992. Collection methods included mist nets placed over bodies of water, capturing bats in residences, and removal of bats from roosts in caves. Several bats were brought to our laboratory by persons who found the bats in their homes.

Individual bats were placed into 250-ml beakers. Each beaker was sealed with a perforated cardboard top, and placed in a refrigerator (4.4°C) for 6-12 h to allow for passage of ingesta that otherwise may have impeded dissection. Bats were euthanized with chloroform and brushed with a toothbrush to collect external parasites for another study (Durden et al., 1992). The gastrointestinal tract (stomach, small and large intestines, and rectum) of each bat was removed, preserved in 10% formalin, and later inspected for parasites with the aid of a dissecting microscope. Parasites discovered in gastrointestinal tracts were removed using a camel-hair brush or dissecting needle and placed into 70% ethanol for storage. Voucher specimens of bats were prepared and deposited in the Auburn University Museum.

One cestode from each infected bat ($n = 14$) and one trematode were examined using scanning electron microscopy. Specimens were dehydrated by placing them into microporous specimen capsules (Electron Microscopy Sciences, Fort Washington, PA), which were placed into vials of 70, 80, 90, and 100% ethanol; 15 min for each change. The final change of ethanol was repeated twice. Alcohol was then removed from dehydrated specimens using a DCP-1 Critical-Point Drying Apparatus (Denton Vacuum, Inc., Cherry Hill, NJ), or by treatment with hexamethyldisilazane (HMDS). Specimens were prepared with HMDS as follows: 100% ethanol was removed from the vial, HMDS was added, and immediately removed; fresh HMDS was added and allowed to remain for 15 min, the HMDS was removed, and fresh HMDS was added for another 15 min; specimens were

then placed onto filter paper in a fume hood to allow for evaporation of the HMDS; fully dried specimens were affixed to carbon tape on aluminum stubs and sputter coated with gold-palladium for 30-50 s. Examination was conducted with a Zeiss DSM 940 digital scanning electron microscope.

Model I contingency table analyses using the G -test of independence were conducted on frequencies of parasitism with respect to taxon of parasite, and species, sex, and collection locality of host. G -test for goodness of fit was used to determine presence or absence of interrelationships in occurrence of parasite taxa (Sokal and Rohlf, 1981).

RESULTS AND DISCUSSION

Of the 324 bats examined, 121 (37.3%) had helminth parasites. Cestodes were found in 14 (4.3%) bats, 47 (14.5%) had nematodes, and 77 (23.8%) had trematodes. Frequency of parasitism and prevalence of parasites for each species of bat are summarized in Table 1. When we compared our results with other surveys of multiple species of bats (Blankespoor and Ulmer, 1970; Nickel and Hansen, 1967; Pistole, 1988), we observed that overall prevalence varied among studies, and that trematodes were the most prevalent helminths, nematodes were next, and cestodes were least prevalent. For example, Nickel and Hansen (1967) surveyed eight species of bats ($n = 65$) from seven counties in Kansas, Nebraska, and Oklahoma, and found that overall prevalence of parasites was 38.5%. When individual taxa were considered, trematodes (26.2%) were most prevalent, followed by nematodes (24.6%) and cestodes (7.7%). Similar results were seen in a study of

Table 1.—Frequencies (F) and prevalences in percent (P) of three taxa of gastrointestinal-helminth parasites in 10 species of bats in Alabama.

Species	n	Cestoidea		Nematoda		Trematoda	
		F	P	F	P	F	P
<i>Eptesicus fuscus</i>	128	0	—	5	3.9	31	24.2
<i>Lasiurus borealis</i>	10	3	30.0	4	40.0	4	40.0
<i>Lasiurus cinereus</i>	1	1	100	1	100	0	—
<i>Lasiurus seminolus</i>	3	3	100	0	—	3	100
<i>Myotis austroriparius</i>	11	2	18.2	2	18.2	6	54.5
<i>Myotis septentrionalis</i>	1	0	—	0	—	1	100
<i>Nycticeius humeralis</i>	15	3	20.0	6	40.0	1	6.7
<i>Pipistrellus subflavus</i>	45	1	2.2	3	6.7	29	64.4
<i>Corynorhinus rafinesquii</i>	1	0	—	0	—	0	—
<i>Tadarida brasiliensis</i>	109	1	0.9	26	23.8	2	1.8

HILTON AND BEST—HELMINTHS OF BATS

64 bats, representing six species from 16 localities in Iowa (Blankespoor and Ulmer, 1970), and in a more recent study of 888 bats, representing nine species, from 65 counties in Indiana (Pistole, 1988).

Only 16 (4.9%) of the parasitized bats in our study had representatives of more than one phylum or class of helminths; three (<1%) were parasitized by nematodes

and trematodes, seven (2.2%) by cestodes and trematodes, five (1.5%) by cestodes and nematodes, and one by cestodes, nematodes, and trematodes (Table 2). Analyses of these data using the *G*-test for goodness of fit showed that bats were most likely to have only cestodes, nematodes, or trematodes, and not a combination of nematode-trematode ($G = 82.89 \gg X^2_{0.05[2]}$), cestode-trematode

Table 2.—Frequencies of Cestoidea (C), Nematoda (N), and Trematoda (T) in bats from Samford Hall, bats from elsewhere in Alabama, and for the two species of bats found in Samford Hall and elsewhere in Alabama.

	<i>n</i>	C	N	T	C and N	C and T	N and T	C, N, and T	Total
Bats from Samford Hall									
<i>Eptesicus fuscus</i>	88	0	2	18	0	0	0	0	20
<i>Tadarida brasiliensis</i>	105	0	25	2	1	0	0	0	28
Total	193	0	27	20	1	0	0	0	48
Bats from elsewhere in Alabama									
<i>Eptesicus fuscus</i>	40	0	1	11	0	0	2	0	14
<i>Lasiurus borealis</i>	10	1	3	1	0	2	1	0	8
<i>Lasiurus cinereus</i>	1	0	0	0	1	0	0	0	1
<i>Lasiurus seminolus</i>	3	0	0	0	0	3	0	0	3
<i>Myotis austroriparius</i>	11	0	2	4	0	2	0	0	8
<i>Myotis septentrionalis</i>	1	0	0	1	0	0	0	0	1
<i>Nycticeius humeralis</i>	15	0	3	0	2	0	0	1	6
<i>Pipistrellus subflavus</i>	45	0	2	29	1	0	0	0	32
<i>Corynorhinus rafinesquii</i>	1	0	0	0	0	0	0	0	0
<i>Tadarida brasiliensis</i>	4	0	0	0	0	0	0	0	0
Total	131	1	11	46	4	7	3	1	73
Bats from Samford Hall and elsewhere in Alabama									
<i>Eptesicus fuscus</i>	128	0	3	29	0	0	2	0	34
<i>Tadarida brasiliensis</i>	109	0	25	2	1	0	0	0	28
Grand Total	324	1	38	66	5	7	3	1	121

($G = 92.17 \gg X^2_{0.05[2]}$), or cestode-nematode ($G = 42.80 \gg X^2_{0.05[2]}$). Individual bats typically host a single taxon of parasite, but dual-taxa parasitizations have been recorded for *L. cinereus* in Maryland (Tromba, 1952) and British Columbia (Webster and Casey, 1973), *N. humeralis* in Indiana (Pistole, 1988), *E. fuscus* in Missouri (Adams and Morris, 1971), and *E. fuscus* and *T. brasiliensis* in Cuba (Barus and del Valle, 1967). Pistole (1988) indicated that *E. fuscus*, *L. borealis*, *M. septentrionalis*, the little brown bat (*Myotis lucifugus*), and the Indiana myotis (*Myotis sodalis*) can be parasitized by more than one taxon of helminth.

That bats often do not harbor parasites from more than one taxon is expected because some bats are dietary specialists (Buchler, 1976; Kunz, 1973). Parasites are obtained from what bats eat (Holmes, 1964; Phillips, 1966) and being a dietary specialist may predispose a bat to become parasitized by only the type of helminth that uses the prey items as intermediate hosts. Although many species of insectivorous bats have a diversified diet that includes several orders and families of insect prey, some species feed selectively on one type of prey and show specificity to foraging habitat. Thus, mayfly (Ephemeroptera) and beetle specialists, such as *M. lucifugus* (Anthony and Kunz, 1977; Buchler, 1976), and moth (Lepidoptera) specialists, such as *L. borealis*, *L. cinereus* (Kunz, 1973; Ross, 1967), and *T. brasiliensis* (Ross, 1967) may be less likely to have dual-taxa parasitizations than *E. fuscus*, which has generalistic feeding habits (Kunz, 1973; Ross, 1967). *L. borealis*, *L. cinereus*, and *M. lucifugus* also exhibit site-specificity of foraging habitats (Kunz, 1973), whereas *E. fuscus* is not site-specific (Furlonger et al., 1987; Geggie and Fenton, 1985; Kunz, 1973).

The only bat in our study that had cestodes, nematodes, and trematodes was a female *N. humeralis* (Table 2) from Fort Rucker, Dale Co. Of parasites of nine adult female and six juvenile *N. humeralis* from Iowa, six adults and one juvenile had cestodes, seven adults and four juveniles had nematodes, and one adult had a trematode (Ubelaker and Kunz, 1971). Two of the taxa must have overlapped and all three may have occurred in one bat. The three taxa do not overlap in any other report of parasites in *N. humeralis* (Alicata, 1932; Chandler, 1938; Chitwood, 1937; Macy and Rausch, 1946; McIntosh, 1932; McIntosh and McIntosh, 1935; Pistole, 1988; Ubelaker, 1966, 1970). Relatively little is known about the biology of *N. humeralis* (Harvey, 1992; Watkins, 1972), but this species may have a

wide range of prey types.

Frequencies of parasitism in *E. fuscus* (22.7%) and *T. brasiliensis* (26.7%) from Samford Hall, Auburn University, were not significantly different ($G = 0.40$, $d.f. = 1$, $P = 0.53$), but frequencies of occurrence of nematodes and trematodes were significantly different between these species ($G = 37.8$, $d.f. = 1$, $P < 0.01$). Of the 20 *E. fuscus* that were parasitized, 10.0% ($n = 2$) had nematodes and 90.0% ($n = 18$) had trematodes (Table 2). Percentages of nematodes and trematodes were opposite for *T. brasiliensis*; 92.8% ($n = 26$) of the 28 parasitized bats had nematodes and 7.1% ($n = 2$) had trematodes (Table 2). Holmes (1964) and Ubelaker (1970) reported that *E. fuscus* is host to more species of trematodes than *T. brasiliensis* and attributed this to differences in feeding habits. Holmes (1964) noted that *T. brasiliensis* feeds heavily on moths and <1% of the diet is composed of insects with aquatic larvae. Conversely, *E. fuscus* primarily feeds on beetles, but >15% of the diet is composed of insects with aquatic larvae. *E. fuscus* also feeds on water scavenger beetles (Coleoptera: Hydrophilidae), mayflies (Ephemeroptera), stoneflies (Plecoptera), caddisflies (Trichoptera), nerve-winged insects (Neuroptera—Hamilton, 1933), predatory diving beetles (Coleoptera: Dytiscidae—Phillips, 1966), diptera flies (Diptera: Dixidae—Ross, 1967), tipulid flies (Diptera: Tipulidae), and midges (Diptera: Chironomidae—Whitaker, 1972). All of these insects have aquatic larvae, and metacercaria of trematodes that infect bats have been found in larvae of midges (McMullen, 1937), caddisflies (Brown, 1933; M. C. Hall, 1929; Knight and Pratt, 1955), stoneflies (J. E. Hall, 1960; M. C. Hall, 1929), and mayflies (Etges, 1959; J. E. Hall, 1960; M. C. Hall, 1929). Conversely, Hamilton (1933) examined 2,200 fecal pellets from *E. fuscus* and found no lepidopteran remains, and Whitaker (1972) found that Lepidoptera accounted for only 4.5% of the diet of 184 *E. fuscus*.

The source of the helminth fauna of *T. brasiliensis* from Samford Hall, Auburn University, is more difficult to explain than that of *E. fuscus*. Life cycles of gastrointestinal nematodes from chiropterans have not been described (Blankespoor and Ulmer, 1970; Ubelaker, 1970), but there are no accounts of nematodes that have lepidopterans as intermediate hosts (Skrjabin et al., 1952, 1954; Yamaguti, 1961). Beetles often serve as intermediate hosts for nematodes parasitic in other mammals (Kinsella, 1991; Morgan and Hawkins, 1951; Ubelaker, 1970). However, *T. brasiliensis* feeds almost exclusively

on moths, and not beetles (Bailey, 1931; Holmes, 1964; Ross, 1967; Storer, 1926). There are at least three possible explanations. First, there may be nematodes that have lepidopterans as intermediate hosts. Second, *T. brasiliensis* is not the moth specialist that the literature indicates. Third, the findings in our study are due to sampling bias; 96.3% ($n = 109$) of *T. brasiliensis* examined by us were collected from one location (Samford Hall, Auburn University). The third possibility may be most likely because *T. brasiliensis* harbors spuriid nematodes (Martin, 1976; Specian and Ubelaker, 1976), which use cockroaches (Blattodea) as intermediate hosts (Specian and Ubelaker, 1976; Ubelaker, 1970). There is a large population of cockroaches on the campus of Auburn University.

Different species of bats from the same collection localities can have dissimilar parasite loads. Font (1978) reported that ca. 80% ($n = 35$) of *M. lucifugus* he examined from Wisconsin harbored a new species of lecithodendriid trematode, *Ototrema schildti*. However, none of the *E. fuscus* ($n = 15$) from the same collection localities were parasitized by *O. schildti*. According to Font (1978), *E. fuscus* is either an unsuitable host for this species of trematode or it does not feed upon insects that have the infective stage. Webster and Casey (1973) found that *E. fuscus* ($n = 24$) and *M. lucifugus* ($n = 31$) from the same three study sites in British Columbia had little overlap in parasites. Only four of the 12 species of helminths that occurred in *E. fuscus* and *M. lucifugus* were found in both species. There were five species of trematodes that occurred in 15 (62.5%) *E. fuscus* compared with three species occurring in six (19.4%) *M. lucifugus*; two species of trematodes were found in both *E. fuscus* and *M. lucifugus*. Nematodes were represented by two species

in 4 (16.7%) *E. fuscus* and three species in 11 (35.5%) *M. lucifugus*; two species of nematodes were shared. One species of cestode was present in one (4.2%) *E. fuscus* and another species was present in two (6.4%) *M. lucifugus* (Webster and Casey, 1973). Webster (1971) examined *T. brasiliensis* ($n = 15$) from a cave in Jamaica and found two species of trematodes in 12 (80.0%) bats and one species of nematode in 9 (60.0%) bats. Parnell's leaf-lipped bats (*Pteronotus parnelli*; $n = 6$) from the same cave did not have trematodes, but was parasitized by three species of nematodes, none of which occurred in *T. brasiliensis*. There also was no overlap in parasites in MacLeay's leaf-lipped bats (*Pteronotus macleayi*; $n = 29$) and *P. parnelli* ($n = 6$) from another cave in Jamaica. These two species of bats failed not only to have overlapping species of parasites, but *P. macleayi* had only trematodes and *P. parnelli* had only nematodes (Webster, 1971).

Of the parasitized bats in our study, 63 (52.1%) were males and 58 (47.9%) were females; frequency of parasitism did not differ by sex ($G = 2.31$, $d.f. = 1$, $P = 0.13$). Sex-specific differences in frequency of parasitism were not apparent for *E. fuscus* ($G = 0.01$, $d.f. = 1$, $P = 0.95$), *P. subflavus* ($G = 1.25$, $d.f. = 1$, $P = 0.26$), or *T. brasiliensis* ($G = 0.68$, $d.f. = 1$, $P = 0.41$; Table 3). Nickel and Hansen (1967) reported that 18 (27.7%) male and 7 (10.8%) female bats were parasitized. Their study included *E. fuscus* ($n = 4$), *P. subflavus* ($n = 10$), and *T. brasiliensis* ($n = 9$), but numbers of males and females of each species were not given. In their study, a male and female *E. fuscus* each had trematodes, one female *P. subflavus* had trematodes, one male *T. brasiliensis* had trematodes, and one female *T. brasiliensis* had nematodes.

Table 3.—Frequencies of parasitism in three species of bats in Alabama based on sex.

Species	Sex	Parasitized	Unparasitized	Total
<i>Eptesicus fuscus</i>	Male	11	31	42
	Female	23	63	86
<i>Pipistrellus subflavus</i>	Male	25	8	33
	Female	7	5	12
<i>Tadarida brasiliensis</i>	Male	12	42	54
	Female	16	39	55
Total		94	188	272

Contingency table analysis of their data failed to reveal a significant difference between these frequencies ($G = 0.03$, $d.f. = 1$, $P = 0.86$). Because of small samples analyzed to date, significant differences between sexes may be detected in future research. Anthony and Kunz (1977) demonstrated that female *M. lucifugus* exhibited a significant preference for beetles and mayflies on a seasonal basis. Unfortunately, no other investigations of parasite faunas included comparisons between sexes for any species of bats, and data from these studies do not allow extraction of such information.

Peak prevalence of parasites for *E. fuscus* from Samford Hall, Auburn University, was in April when 40.0% ($n = 5$) were hosts to helminths. Prevalence declined monthly thereafter until it reached 8.3% ($n = 12$) in October (Fig. 1a). Prevalences for *T. brasiliensis* from Samford Hall had a more irregular distribution. Greatest prevalence was in November when 50.0% ($n = 12$) had parasites and lowest prevalence was in September when none of the bats ($n = 12$) were parasitized (Fig. 1b).

Coggins et al. (1982) found that *M. lucifugus* in Wisconsin had lowest prevalences of parasites in summer (July and August) and highest prevalences in spring (April, May, and June) and autumn (September and October). However, Blankespoor and Ulmer (1970) and Nickel and Hansen (1967) found that prevalences were low in spring and increased until the bats entered hibernation in autumn. The authors of both studies suggested that bats lost their parasites during hibernation and were reinvaded by parasites when they came out of hibernation (Blankespoor and Ulmer, 1970; Nickel and Hansen, 1967). According to Coggins et al. (1982), this strategy would work only for parasites of bats that did not undergo long (>3 months) periods of hibernation. Because bats in Wisconsin hibernate for nearly 8 months, the months of activity are not long enough to ensure successful completion of life cycles of parasites (Coggins et al., 1982). Coggins et al. (1982) maintain that parasites of bats from northerly latitudes are adapted to overwintering in their definitive host and not the intermediate host as do most helminths. One would expect prevalences in *E. fuscus* and *T. brasiliensis* in Alabama to follow the scenario proposed by Blankespoor and Ulmer (1970) and Nickel and Hansen (1967). Instead, prevalences in *E. fuscus* were highest in spring and lowest in autumn, and prevalences in *T. brasiliensis* failed to follow a discernable pattern.

Scanning electron micrographs were taken of at least one cestode from every bat that harbored cestodes; a




Fig. 1.—Number of a) *Eptesicus fuscus* and b) *Tadarida brasiliensis* from Samford Hall, Auburn University, Lee Co., Alabama, examined and prevalence of parasites by month during 1990.

trematode from one *L. borealis* also was micrographed. Cestodes from *L. seminolus* did not appear to have a rostellum or armed suckers, which suggests that they belong to the Anoplocephalidae (Schmidt, 1986). It was not apparent whether the cestode from *M. austroriparius* had a rostellum or whether the suckers were armed. The cestode from *P. subflavus* appeared to have a rostellum and unarmed suckers, which suggested that it was a member of the Hymenolepididae (Schmidt, 1986). The trematode from *L. borealis* appeared to be a distome, because it had an oral sucker and an acetabulum on the mid-ventral surface (Schmidt and Roberts, 1989). Measurements were not taken of potentially diagnostic features because parasites were fixed in situ. This fixation method causes a layer of mucus to become affixed to specimens, often distorts specimens by causing them to contract, and may result in diagnostic features such as armature becoming

invaginated (M. H. Pritchard, pers. comm.; Pritchard and Kruse, 1982). Consequently, diagnostic features were not visible or were distorted such that there was a significant probability of reaching erroneous taxonomic conclusions. However, Hymenolepididae and Anoplocephalidae are the only families of cestodes to be reported in bats from the United States (Ubelaker, 1970), and the trematodes of bats in the United States are distomes (Schmidt and Roberts, 1989; Yamaguti, 1958).

ACKNOWLEDGMENTS

This paper was extracted from a graduate thesis (Hilton, 1994) that was submitted in partial fulfillment of requirements for the Master of Science degree in Zoology at Auburn University. We thank R. S. Boyd for assistance with statistical analyses, R. R. Dute for help with electron microscopy and micrography, the Alabama Department of Conservation and Natural Resources for providing a scientific-collecting permit, and D. Bond, S. L. Burt, T. H. Haas, J. L. Hilton, K. Youngblood, and other persons that helped with field or laboratory work. G. R. Hepp, C. A. Sundermann, and L. C. Wit critically evaluated an early draft of the manuscript. This is journal article 15-944833 of the Alabama Agricultural Experiment Station.

LITERATURE CITED

- Adams, G. J., and T. A. Morris. 1971. A preliminary investigation of the parasites of certain hibernating bats common to southwestern Missouri. *Transactions of the Missouri Academy of Science*, 5:122.
- Alicata, J. E. 1932. A new trematode of the genus *Urotrema* from bats. *Proceedings of the United States National Museum*, 81:1-4.
- Anthony, E. L. P., and T. H. Kunz. 1977. Feeding strategies of the little brown bat, *Myotis lucifugus*, in southern New Hampshire. *Ecology*, 58:775-786.
- Bailey, V. 1931. *Mammals of New Mexico*. North American Fauna, 53:1-412.
- Barus, V., and M. T. del Valle. 1967. Systematic survey of nematodes parasitizing bats (Chiroptera) in Cuba. *Folia Parasitologica (Praha)*, 14:121-140.
- Best, T. L., S. D. Carey, K. G. Caesar, and T. H. Henry. 1993. Distribution and abundance of bats (Mammalia: Chiroptera) in Coastal Plain caves of southern Alabama. *National Speleological Society Bulletin*, 54:61-65.
- Blankespoor, H. D., and M. J. Ulmer. 1970. Helminths from six species of Iowa bats. *Proceedings of the Iowa Academy of Science*, 77:200-206.
- . 1972. *Prosthodendrium volaticum* sp. n. (Trematoda: Lecithodendriidae) from two species of Iowa bats. *Proceedings of the Helminthological Society of Washington*, 39:224-226.
- Bray, R. S. 1984. Some parasitic protozoa from the Gambia. *Journal of Protozoology*, 31:577-578.
- Brown, F. J. 1933. On the excretory system and life history of *Lecithodendrium chilostromum* (Mehl.) and other bat trematodes, with a note on the life history of *Dicrocoelium dendriticum* Rudolphi. *Parasitology*, 25:317-329.
- Buchler, E. 1976. Prey selection by *Myotis lucifugus* (Chiroptera: Vespertilionidae). *The American Naturalist*, 110:619-628.
- Byrd, E. E., and R. W. Macy. 1942. Mammalian trematodes. III. Certain species from bats. *Journal of the Tennessee Academy of Science*, 17:149-156.
- Chandler, A. C. 1938. A report on the parasites of a bat, *Nycticeius humeralis*, with descriptions of four new helminths. Pp. 107-114, in *Livro jubilar Professor Lauro Travassos* (A. Neiva, ed.). Instituto Oswaldo Cruz, Rio de Janeiro, Brazil, 294 pp.
- Chitwood, B. G. 1937. A new trichostrongyle, *Allintoshius nycticeius* n. g., n. sp. (Nematoda) from a bat. *Proceedings of the Helminthological Society of Washington*, 4:19-20.
- Coggins, J. R. 1988. Methods for the ecological study of bat endoparasites. Pp. 475-489, in *Ecological and behavioral methods for the study of bats* (T. H. Kunz, ed.). Smithsonian Institution Press, Washington, D.C., 533 pp.
- Coggins, J. R., J. L. Tedesco, and C. Rupprecht. 1982. Seasonal changes and overwintering of parasites in the bat, *Myotis lucifugus* (Le Conte), in a Wisconsin hibernaculum. *The American Midland Naturalist*, 107:305-315.
- Durden, L. A., T. L. Best, N. Wilson, and C. D. Hilton. 1992. Ectoparasitic mites (Acari) of sympatric Brazilian free-tailed bats and big brown bats in Alabama. *Journal of Medical Entomology*, 29:507-511.
- Duszynski, D. W., D. W. Reduker, and B. P. Parker. 1988. *Eimeria* from bats of the world. II. A new species in *Tadarida femorasacca* from Sonora, Mexico. *The Journal of Parasitology*, 74:317-321.
- Etges, F. J. 1959. Studies on the life history of a species of *Acanthatrium* (Trematoda: Lecithodendriidae).

- The Journal of Parasitology, 45 (supplement):18.
- Fischthal, J. H., and R. L. Martin. 1978. *Postorchigenes paraguayensis* sp. n. (Trematoda, Pleurogenidae), a digenetic trematode from a large fishing bat, *Noctilio leporinus rufescens* Olfers, from Paraguay. Acta Parasitologica Polonica, 25:217-221.
- Font, W. F. 1978. *Ototrema schildti* gen. et sp. n. (Trematoda: Lecithodendriidae) from the little brown bat, *Myotis lucifugus*, in Wisconsin. The Journal of Parasitology, 64:391-392.
- Foster, G. W. 1979. *Polychromophilus* from southeastern brown bats (*Myotis austroriparius*) in north-central Florida. The Journal of Parasitology, 65:465-466.
- Furlonger, C. L., H. J. Dewar, and M. B. Fenton. 1987. Habitat use by foraging insectivorous bats. Canadian Journal of Zoology, 65:284-288.
- Geggie, J. F., and M. B. Fenton. 1985. A comparison of foraging by *Eptesicus fuscus* (Chiroptera: Vespertilionidae) in urban and rural environments. Canadian Journal of Zoology, 63:263-267.
- Greenhall, A. M., and U. Schmidt. 1988. Natural history of vampire bats. CRC Press, Boca Raton, Florida, 246 pp.
- Hall, E. R. 1981. The mammals of North America. Second ed. John Wiley & Sons, New York, 1:1-600 + 90.
- Hall, J. E. 1960. Some lecithodendriid metacercaria from Indiana and Michigan. The Journal of Parasitology, 46:309-315.
- Hall, M. C. 1929. Arthropods as intermediate hosts of helminths. Smithsonian Miscellaneous Collections, 81:1-77.
- Hamilton, W. J., Jr. 1933. The insect food of the big brown bat. Journal of Mammalogy, 14:155-156.
- Harvey, M. J. 1992. Bats of the eastern United States. Arkansas Game and Fish Commission, Little Rock, 46 pp.
- Henry, T. H., T. L. Best, and C. D. Hilton. 2000. Body size, reproductive biology, and sex ratio of a year-round colony of *Eptesicus fuscus fuscus* and *Tadarida brasiliensis cynocephala* in eastern Alabama. Occasional Papers of the North Carolina Museum of Natural Sciences and the North Carolina Biological Survey, 12:50-66.
- Hilton, C. D. 1994. Gastrointestinal helminth parasites of bats (Chiroptera: Molossidae, Vespertilionidae) in Alabama. M.S. thesis, Auburn University, Alabama, 65 pp.
- Holmes, J. C. 1964. Factors influencing the trematode fauna of bats. Proceedings of the First International Congress on Parasitology, 1:490-492.
- Kinsella, J. M. 1991. Comparison of helminths of three species of mice, *Peromyscus floridanus*, *Peromyscus gossypinus*, and *Peromyscus polionotus*, from southern Florida. Canadian Journal of Zoology, 69:3078-3083.
- Knight, R. A., and I. Pratt. 1955. The life histories of *Allassogonoporus vespertilionis* Macy and *Acanthatrium oregonense* Macy (Trematoda: Leuchodendriidae). The Journal of Parasitology, 41:248-255.
- Kunz, T. H. 1973. Resource utilization: temporal and spatial components of bat activity in central Iowa. Journal of Mammalogy, 54:14-32.
- Loftin, H. 1961. An annotated check-list of trematodes and cestodes and their vertebrate hosts from Northwest Florida. Quarterly Journal of the Florida Academy of Sciences, 23:302-314.
- Macy, R. W., and R. L. Rausch. 1946. Morphology of a new species of bat cestode, *Hymenolepis roudabushi*, and a note on *Hymenolepis christensoni*, Macy. Transactions of the American Microscopical Society, 65:173-175.
- Martin, D. R. 1976. New host and distribution records of helminth parasites of the Brazilian free-tail bat, *Tadarida brasiliensis*, from Texas and Louisiana. Proceedings of the Helminthological Society of Washington, 43:85-86.
- McIntosh, A. 1932. A new filarid from the abdominal cavity of a bat. The Journal of Parasitology, 19:170.
- McIntosh, A., and G. E. McIntosh. 1935. Additional notes on two bat parasites, *Dicrocoelium lasiuri* McIntosh, 1933 (Trematoda: Dicrocoeliidae) and *Litomosa americana* McIntosh, 1932 (Nematoda: Filariidae). Proceedings of the Helminthological Society of Washington, 2:60-62.
- McMullen, D. B. 1937. The life histories of three trematodes, parasitism in birds and mammals, belonging to the genus *Plagiorchis*. The Journal of Parasitology, 23:235-243.
- Morgan, B. B., and P. A. Hawkins. 1951. Veterinary helminthology. Burgess Publishing Company, Minneapolis, Minnesota, 400 pp.
- Nickel, P. A., and M. F. Hansen. 1967. Helminths of bats collected in Kansas, Nebraska and Oklahoma. The American Midland Naturalist, 78:481-486.

- Peterson, R. L., and P. Kirmse. 1969. Notes on *Vampyrum spectrum*, the false vampire bat. *Canadian Journal of Zoology*, 47:140-142.
- Phillips, G. L. 1966. Ecology of the big brown bat (Chiroptera: Vespertilionidae) in northeastern Kansas. *The American Midland Naturalist*, 75:168-198.
- Pistole, D. H. 1988. A survey of helminth parasites of chiropterans from Indiana. *Proceedings of the Helminthological Society of Washington*, 55:270-274.
- Pritchard, M. H., and G. O. W. Kruse. 1982. The collection and preservation of animal parasites. University of Nebraska Press, Lincoln, 141 pp.
- Ross, A. 1967. Ecological aspects of the food habits of insectivorous bats. *Proceedings of the Western Foundation of Vertebrate Zoology*, 1:204-263.
- Schmidt, G. D. 1986. Handbook of tapeworm identification. CRC Press, Boca Raton, Florida, 675 pp.
- Schmidt, G. D., and L. S. Roberts. 1989. Foundations of parasitology. Fourth ed. Times Mirror/Mosby College Publishing, St. Louis, Missouri, 750 pp.
- Skrjabin, K. I., N. P. Shikhobalova, and R. S. Schulz. 1954. Essentials of nematology. Trichostrongylids of animals and man. Israel Program for Scientific Translations, Jerusalem, Israel, and the United States Department of Agriculture, Washington, D.C., 3:1-704.
- Skrjabin, K. I., N. P. Shikhobalova, R. S. Schulz, T. I. Popova, S. N. Boev, and S. L. Delyamure. 1952. Key to parasitic nematodes. Strongylata. Israel Program for Scientific Translations, Jerusalem, Israel, and the United States Department of Agriculture, Washington, D.C., 3:1-890.
- Sokal, R. R., and F. J. Rohlf. 1981. Biometry: the principles and practice of statistics in biological research. Second ed. W. H. Freeman and Company, San Francisco, 859 pp.
- Specian, R. D., and J. E. Ubelaker. 1976. Redescription of a nematode, *Seuratium cancellatum* Chitwood, 1938, from bats in Texas. *Proceedings of the Helminthological Society of Washington*, 43:59-65.
- Stiles, C. W., and M. O. Nolan. 1931. Key-catalogue of parasites reported for Chiroptera (bats) with their possible public health importance. *National Institutes of Health Bulletin*, United States Treasury Department, 155:603-742.
- Storer, T. I. 1926. Bats, bat towers and mosquitos. *Journal of Mammalogy*, 7:85-90.
- Tromba, F. G. 1952. *Myotis lucifugus*; a new aberrant host for the third stage larva of *Physocephalus sexalatus* (Molin, 1860). *The Journal of Parasitology*, 38:497.
- Ubelaker, J. E. 1966. Parasites of the gray bat, *Myotis grisescens*, in Kansas. *The American Midland Naturalist*, 75:199-204.
- . 1970. Some observations on ecto- and endoparasites of Chiroptera. Pp. 247-261, in *About bats: a chiropteran biology symposium* (B. H. Slaughter and D. W. Walton, eds.). Southern Methodist University Press, Dallas, Texas, 339 pp.
- Ubelaker, J. E., and T. H. Kunz. 1971. Parasites of the evening bat, *Nycticeius humeralis*, in Iowa. *The Texas Journal of Science*, 22:425-427.
- Ubelaker, J. E., R. D. Specian, and D. W. Duszynski. 1977. Endoparasites. Pp. 7-56, in *Biology of bats of the New World family Phyllostomatidae* (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Special Publications, The Museum, Texas Tech University, 13:1-364.
- Watkins, L. C. 1972. *Nycticeius humeralis*. *Mammalian Species*, 23:1-4.
- Webster, W. A. 1971. Studies on the parasites of Chiroptera. I. Helminths of Jamaican bats of the genera *Tadarida*, *Chilonycteris*, and *Monophyllus*. *Proceedings of the Helminthological Society of Washington*, 38:195-199.
- . 1973. Studies on the parasites of Chiroptera. II. Checklist of the helminth parasites of North American bats. *Wildlife Disease*, 59:1-26.
- Webster, W. A., and G. A. Casey. 1973. Studies on the parasites of Chiroptera. III. Helminths from various bat species collected in British Columbia. *Canadian Journal of Zoology*, 51:633-636.
- Wheat, B. E. 1975. *Eimeria macyi* sp. n. (Protozoa: Eimeridae) from the eastern pipistrelle, *Pipistrellus subflavus* from Alabama. *The Journal of Parasitology*, 61:920-922.
- Whitaker, J. O., Jr. 1972. Food habits of bats from Indiana. *Canadian Journal of Zoology*, 50:877-883.
- White, J. S. 1959. The acarine ectoparasites of Alabama bats. Ph.D. dissertation, University of Alabama, Tuscaloosa, 137 pp.
- Yamaguti, S. 1958. *Systema Helminthum: the digenetic trematodes of vertebrates*. Interscience Publishers, Inc., John Wiley & Sons, New York, 1:1-1575.
- . 1961. *Systema Helminthum: the nematodes of vertebrates*. Interscience Publishers, Inc., John Wiley & Sons, New York, 3:1-679.

Zdzitowiecki, K., and M. A. Rutkowska. 1980. The helminthofauna of bats (Chiroptera) from Cuba. III. A review of trematodes. *Acta Parasitologica Polonica*, 26:201-214.

APPENDIX I

Specimens Examined

Eptesicus fuscus—*Butler Co.*: Georgiana, bridge on state road 106 where it passes over West Railroad Avenue (mile marker 12) (6 males, 19 females). *Lee Co.*: Auburn (0, 1); Auburn University, attic of Samford Hall (28, 60); Opelika, 1103 Collinwood Drive (5, 6); Opelika, inside duplex at 709 North 8th Street (1, 0). *Morgan Co.*: Talucah, Talucah Cave (2, 0).

Lasiurus borealis—*Dale Co.*: Fort Rucker, Ech Lake (0, 3); Fort Rucker, Girl Scout Camp (1, 3). *Lee Co.*: Auburn University, Ralph B. Draughon Library (1, 0); Flake Creek at county road 65, 0.3 mile from state road 169 (0, 1); Pond on state road 169, 0.3 mile from county road 65, teardrop-shaped pond (0, 1).

Lasiurus cinereus—*Lee Co.*: Farm pond 0.98 mile from state road 169 on county road 81 (1, 0).

Lasiurus seminolus—*Dale Co.*: Fort Rucker, Ech Lake (0, 1); Fort Rucker, Girl Scout Camp (0, 1); Fort Rucker, headquarters (1, 0).

Myotis austroriparius—*Conecuh Co.*: Hodges Cave (1, 0); Sanders Cave, 3 miles NW Brooklyn (2, 0). *Covington Co.*: Rock House Cave, 7 miles NE Florala (1, 0). *Monroe Co.*: Locklin Cave, T6N R5E Section 4 (4, 0); Locklin Cave, 5 miles W Perdue Hill (3, 0).

Myotis septentrionalis—*Franklin Co.*: 9 miles NE Red Bay on county road 88 before the Bear Creek Levee (1, 0).

Nycticeius humeralis—*Dale Co.*: Fort Rucker, Girl Scout Camp (3, 2). *Lee Co.*: Flake Creek at county road 65, 0.3 mile from state road 169 (1, 0); Opelika, 209 South 4th Street (Bethesda Baptist Church) (0, 3); Opelika, 403 Avenue A (0, 2); Opelika, 403 Avenue A and attic of Bethesda Baptist Church across street from 403 Avenue A (1, 3).

Pipistrellus subflavus—*Butler Co.*: B. C. Barganier Cave (=Rock Cave) (3, 0). *Clark Co.*: Buzzard's Den Cave, T9N R1E Section 18 (11, 0); Lion's Den Cave, 3 miles SW McEntyre (1, 0). *Conecuh Co.*: Sanders Cave, 3 miles NW Brooklyn (3, 2). *Franklin Co.*: 9 miles NE Red Bay on county road 88 before the Bear Creek Levee (5, 4); Goat Cave (=Belgreen Underground Lake), near Belgreen (2, 1). *Lee Co.*: Flake Creek at county road 65, 0.3 mile from state road 169 (1, 0); Pond on state road 169, 0.3 mile from county road 65, teardrop-shaped pond (1, 0). *Marshall Co.*: Mike's Wolff Cave, Union Grove (4, 0). *Marshall Co.*: Wolff Cave, Union Grove (0, 5). *Morgan Co.*: Talucah Cave, Talucah (2, 0).

Corynorhinus rafinesquii—*Clark Co.*: Buzzard's Den Cave, T9N R1E Section 18 (1, 0).

Tadarida brasiliensis—*Bullock Co.*: Union Springs, 304 East Hardaway (0, 1). *Butler Co.*: Georgiana, bridge on state road 106 where it passes over West Railroad Avenue (mile marker 12) (2, 0). *Lee Co.*: Auburn University, attic of Samford Hall (51, 54); Opelika, 1103 Collinwood Drive (1, 0).